



## The Co-Transcriptome of Uropathogenic Escherichia coli-Infected Mouse Macrophages Reveals New Insights into Host-Pathogen Interactions

Journal:	<i>Cellular Microbiology</i>
Manuscript ID:	CMI-14-0241.R1
Manuscript Type:	Research article
Date Submitted by the Author:	01-Nov-2014
Complete List of Authors:	<p>Mavromatis, Charalampos; KAUST, Division of Biological and Environmental Sciences &amp; Engineering            Bokil, Nilesh; The University of Queensland, IMB            Totsika, Makrina; The University of Queensland, IMB            Kakkanat, Asha; University of Queensland, Australian Infectious Diseases Research Centre            Schaale, Kolja; The University of Queensland, IMB            Cannistraci, Carlo; Technische Universität Dresden, Biotec            Ryu, Taewoo; KAUST, Division of Biological and Environmental Sciences &amp; Engineering            Beatson, Scott; University of Queensland, School of Chemistry and Molecular Biosciences and Australian Infectious Diseases Research Centre            Ulett, Glen; Griffith University, School of Medical Sciences            Schembri, Mark; University of Queensland, School of Molecular and Microbial Sciences            Sweet, Matthew J.; The University of Queensland, IMB            Ravasi, Timothy; King Abdullah University of Science and Technology, Division of Biological and Environmental Sciences &amp; Engineering</p>
Key Words:	Genomics, Immunology, Infection, Mechanism of action, Bioinformatics, Transcriptomics

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**The Co-Transcriptome of Uropathogenic *Escherichia coli*-Infected Mouse Macrophages  
Reveals New Insights into Host–Pathogen Interactions**

Charalampos (Harris) Mavromatis<sup>1,2</sup>, Nilesh J. Bokil<sup>3,4</sup>, Makrina Totsika<sup>4,5,6</sup>, Asha Kakkanat<sup>4,5</sup>, Kolja  
Schaale<sup>3,4</sup>, Carlo V. Cannistraci<sup>1,2,7</sup>, Taewoo Ryu<sup>1,2</sup>, Scott A. Beatson<sup>4,5</sup>, Glen C Ulett<sup>8</sup>, Mark A.  
Schembri<sup>4,5</sup>, Matthew J. Sweet<sup>3,4</sup> and Timothy Ravasi<sup>1,2§</sup>

1. Division of Biological and Environmental Sciences and Engineering, Division of Computer, Electrical and Mathematical Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal, Kingdom of Saudi Arabia
2. Division of Medical Genetics, Department of Medicine, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA USA
3. Institute for Molecular Bioscience, The University of Queensland, St Lucia, Australia
4. Australian Infectious Diseases Research Centre, The University of Queensland, St Lucia, Australia
5. School of Chemistry and Molecular Biosciences, The University of Queensland, St Lucia, Australia
6. Institute of Health and Biomedical Innovation (IHBI), School of Biomedical Sciences, Queensland University of Technology (QUT), 60 Musk Ave/cnr. Blamey St, Kelvin Grove, QLD, 4059, Australia
7. Biomedical Cybernetics Group, Biotechnology Center (BIOTEC), Technische Universität Dresden, Tatzberg 47/49, 01307 Dresden, Germany.
8. Griffith Health Institute and School of Medical Science, Griffith Health Centre, Gold Coast Campus, Griffith University, Southport, QLD 4222, Australia.

§ Corresponding author

**Email Addresses of Authors**

CM <harris.mavromatis@kaust.edu.sa>

NJB <n.bokil@imb.uq.edu.au>

MT < makrina.totsika@qut.edu.au >

AK <asha.kakkanat@uq.net.au>

KS <k.schaale@imb.uq.edu.au>

CVC <carlo.cannistraci@biotec.tu-dresden.de>

TR <taewoo.ryu@kaust.edu.sa>

SAB <s.beatson@uq.edu.au>

GCC <g.ulett@griffith.edu.au>

MAS <m.schembri@uq.edu.au>

MJS <m.sweet@imb.uq.edu.au>

TR <timothy.ravasi@kaust.edu.sa>

**Word Count: 6,735**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Summary**

Urinary tract infections (UTI) are among the most common infections in humans. Uropathogenic *Escherichia coli* (UPEC) can invade and replicate within bladder epithelial cells, and some UPEC strains can also survive within macrophages. To understand the UPEC transcriptional program associated with intramacrophage survival, we performed host–pathogen co-transcriptome analyses using RNA sequencing. Mouse bone marrow-derived macrophages (BMMs) were challenged over a 24 h time course with two UPEC reference strains that possess contrasting intramacrophage phenotypes: UTI89, which survives in BMMs, and 83972, which is killed by BMMs. Neither of these strains caused significant BMM cell death at the low multiplicity of infection that was used in this study. We developed an effective computational framework that simultaneously separated, annotated, and quantified the mammalian and bacterial transcriptomes. BMMs responded to the two UPEC strains with a broadly similar gene expression program. In contrast, the transcriptional responses of the UPEC strains diverged markedly from each other. We identified UTI89 genes upregulated at 24 h post-infection, and hypothesized that some may contribute to intramacrophage survival. Indeed, we showed that deletion of one such gene (*pspA*) significantly reduced UTI89 survival within BMMs. Our study provides a technological framework for simultaneously capturing global changes at the transcriptional level in co-cultures, and has generated new insights into the mechanisms that UPEC use to persist within the intramacrophage environment.

## Introduction

Urinary tract infections (UTIs) represent one of the most significant community-acquired and healthcare-associated diseases (Foxman, 2010, Horvath *et al.*, 2012). Uncomplicated UTIs result in more than 14 million medical visits and account for almost \$4 billion in medical expenditure each year in the USA alone (Salvatore *et al.*, 2011). Approximately 50% of women will experience a UTI at some point in their life, with almost 25% of patients experiencing a recurrence within the first 6 months following treatment of the initial UTI (Salvatore *et al.*, 2011). An estimated 68% of recurrent UTIs arise from the same bacterial strain that caused the initial infection (Hunstad *et al.*, 2010). Uropathogenic *Escherichia coli* (UPEC) is the most common causative agent of UTIs, being responsible for ~80% of all community-acquired infections (Foxman, 2010). In the majority of acute, uncomplicated UPEC-mediated UTIs, single cultured isolates are diagnostic of the infection (Willner *et al.*, 2014).

UPEC employ a range of virulence factors, including adhesins, toxins and iron-acquisition systems, to colonize the urinary tract and cause disease (Totsika *et al.*, 2012, Ulett *et al.*, 2013). Different UPEC strains display extensive genetic diversity owing to the presence of mobile DNA elements such as “pathogenicity islands”, prophages and plasmids (Hacker *et al.*, 2000, Mysorekar *et al.*, 2006, Wiles *et al.*, 2008, Hunstad *et al.*, 2010, Hannan *et al.*, 2012). The UPEC strains UTI89 (Mulvey *et al.*, 2001) and 83972 (Lindberg *et al.*, 1975c, Klemm *et al.*, 2007, Zdziarski *et al.*, 2010) are representative of cystitis and asymptomatic bacteriuria (ABU) isolates, respectively.

Acute pyelonephritis and ABU represent the two extremes of UTI. Acute pyelonephritis is a severe, acute systemic infection caused by UPEC strains containing virulence genes clustered on pathogenicity islands (Eden *et al.*, 1976, Funfstuck *et al.*, 1986, Stenqvist *et al.*, 1987, Orskov *et al.*, 1988, Johnson, 1991, Welch *et al.*, 2002). ABU, on the other hand, is an asymptomatic carrier state that resembles commensalism. A single *E. coli* strain may be present in ABU patients at levels of more

1  
2 than  $10^5$  colony-forming units (CFU)  $\text{ml}^{-1}$  for months or years without provoking a host response.  
3  
4 Because the majority of ABU-associated *E. coli* strains are non-hemolytic, non-adherent and lack  
5  
6 hemagglutination ability, early studies suggested that this behavior reflected a lack of virulence genes  
7  
8 (Lindberg, 1975, Lindberg *et al.*, 1975a, Lindberg *et al.*, 1975b, Lindberg *et al.*, 1975c, Eden *et al.*,  
9  
10 1976, Kaijser *et al.*, 1977). Molecular epidemiology has shown, however, that many ABU strains carry  
11  
12 virulence genes despite failing to express the phenotype (Plos *et al.*, 1990, Plos *et al.*, 1995, Mabbett  
13  
14 *et al.*, 2009).

15  
16  
17 As with all infectious agents, UPEC must overcome innate immunity, a biological system  
18  
19 compromising both cellular mediators (e.g. neutrophils) and soluble mediators (e.g. complement  
20  
21 proteins) that act synergistically. Several studies have investigated the role of neutrophils in UPEC-  
22  
23 mediated pathology (Ingersoll *et al.*, 2008, Sivick *et al.*, 2010, Lau *et al.*, 2012, Tourneur *et al.*, 2013),  
24  
25 whereas there is a paucity of information on the interactions between UPEC and macrophages,  
26  
27 another key cellular component of innate immunity (Tegner *et al.*, 2006). We previously demonstrated  
28  
29 that the ability of UPEC to survive in mouse macrophages differs markedly between different strains;  
30  
31 some strains, such as UTI89, are able to survive over a 24-hour infection period, whereas others, such  
32  
33 as 83972, are rapidly killed (Bokil *et al.*, 2011). This suggests that UPEC strains like UTI89 are able to  
34  
35 subvert macrophage antimicrobial pathways, though the mechanisms responsible are still unknown.  
36  
37 These findings are in keeping with a larger body of literature documenting intraepithelial cell survival of  
38  
39 some UPEC strains, both *in vitro* and *in vivo* (Hunstad *et al.*, 2010).  
40  
41  
42

43 Next-generation sequencing technologies provide a powerful approach for studying co-  
44  
45 transcriptomics during infection (t Hoen *et al.*, 2008, Hegedus *et al.*, 2009, Jager *et al.*, 2009, Xiang *et*  
46  
47 *al.*, 2010b, Huang *et al.*, 2012, Nie *et al.*, 2012, Wang *et al.*, 2012). These direct sequencing  
48  
49 methodologies allow the measurement of millions of RNA transcripts in a sample, thus enabling  
50  
51 identification of global differences in gene expression under different growth conditions (Morozova *et*  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 *al.*, 2008, Wang *et al.*, 2009). RNA sequencing generates information about absolute transcript levels,  
3  
4 avoiding many of the limitations of microarrays (t Hoen *et al.*, 2008, Llorens *et al.*, 2011). To further  
5  
6 understand the transcriptional programs that are simultaneously activated during host-pathogen  
7  
8 interaction, we developed an approach for isolating total RNA from co-cultures and analyzing the  
9  
10 simultaneous changes in expression that take place in the interacting organisms. Previous studies  
11  
12 (Hegedus *et al.*, 2009, Xiang *et al.*, 2010a, Xiao *et al.*, 2010, Ordas *et al.*, 2011) have focused on  
13  
14 either the host or the pathogen, without revealing simultaneous co-transcriptomic changes that occur  
15  
16 during infection. Such methods have also relied on the isolation of species-specific RNA, which can  
17  
18 introduce biases in the analysis.  
19  
20

21 To gain insights into novel strategies used by UPEC to subvert macrophage anti-microbial  
22  
23 responses, we performed a global co-transcriptomic analysis of UPEC gene expression within murine  
24  
25 bone marrow-derived macrophages (BMMs). We used RNA sequencing to monitor (i) the  
26  
27 transcriptional responses of UPEC strains UTI89 and 83972 within the intramacrophage environment  
28  
29 across an extended time course, and (ii) differences in macrophage gene-expression responses to  
30  
31 each strain. Our comprehensive approach has generated new insights into host–pathogen interactions  
32  
33 and the possible consequences of these interactions for disease processes.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Results**

*Analysis of Digital Gene Expression Libraries*

The UPEC strains UTI89 and 83972 display contrasting intramacrophage survival phenotypes; UTI89 is able to survive in significant numbers, whereas 83972 is rapidly killed (Bokil *et al.*, 2011). To examine the molecular basis for this difference, we investigated the gene expression profiles of UTI89 and 83972 in BMMs in parallel over a 24 h period. For our experimental system, we used a multiplicity of infection (MOI) of 10:1, since this MOI does not have obvious effects on BMM cell viability for either of the strains. As expected, bacterial loads of UTI89 that were recovered from BMM were substantially higher than those of 83972 (Supplementary Figure 1). Total RNA was harvested at 2, 4 and 24 h post-infection (hpi) and global gene expression profiles were analyzed using the Illumina Hi-Seq 2000 Digital Gene Expression Tag Profiling Kit, a tag-based transcriptome sequencing method. cDNA libraries were prepared from gentamicin-treated UPEC-BMM co-cultures, sequenced and analyzed together with bacterial and BMM control samples. The RNA-Seq files generated for all libraries were preprocessed by a custom java script (Supplementary Table 1).

*Mapping RNA-Seq Reads*

Alignment of sequencing reads to the respective mouse and UPEC reference genomes was performed using TopHat. Sequences were mapped against their respective *Mus musculus* or *E. coli* genomes. The resulting BAM files were further used to compute alignment statistics for all libraries employing the samtool flagstat command. These processes are summarized in Supplementary Table 2. As expected, very few sequence reads were captured from 83972-BMM co-cultures at 24 hpi, consistent with the observation that these bacteria were essentially cleared by BMMs at this time point. Hence, subsequent analyses of gene expression in 83972 excluded this specific condition.



### BMM Transcriptional Responses to UPEC

To explore transcriptional relationships between different conditions, we performed dimensionality reduction analysis using minimum curvilinearity embedding (MCE) (Cannistraci *et al.*, 2010, Cannistraci *et al.*, 2013). This analysis revealed that the BMM genes whose expression was regulated by infection were not markedly different between the two different UPEC strains at the initial stage, 2hpi. However, after 4hpi, a slight difference emerged, which was also maintained at 24hpi. As expected, however, we observed that the time-dependent regulation of gene expression (Figure 1A) was the main pattern that emerged from the data, indicating that the macrophage transcriptional response follows a distinct temporal profile that is common to infection by both strains, and much more distinctive than the difference between the two strains.

To gain further insights into the global transcriptional changes that take place in UPEC-infected macrophages, we catalogued differentially expressed genes (DEG) in normalized digital gene expression data through pairwise comparisons between controls (uninfected BMM cultures) and treatments (co-cultures of UPEC-infected BMMs) using a previously described method (Trapnell *et al.*, 2012) with a threshold of a false-discovery rate (FDR)-adjusted  $P$ -value  $< 0.01$  in at least one of the pairwise comparisons. Using this approach, we identified 628 and 652 BMM genes that were differentially regulated following infection with UTI89 or 83972, respectively, over a 24-hour infection time course. Further analyses of the BMM transcriptome identified 603 genes that were commonly regulated by UTI89 and 83972, as well as 25 and 49 BMM genes that were differentially regulated after infection with UTI89 or 83972, respectively (Figure 1B). The greatest divergence in responses to each strain occurred at the latest time point post-infection.

To investigate the regulatory patterns of divergently expressed (DE) genes, we clustered genes as either up or downregulated. This analysis revealed a rapid response for upregulated genes

1  
2 that was maintained throughout the infection time course. In contrast, the number of downregulated  
3  
4 genes was much lower at 2 and 4 hpi, but increased markedly by 24 hpi (Figure 1C). The substantial  
5  
6 overlap in DEG, as well as their conserved pattern of regulation, suggests that the macrophage  
7  
8 transcriptional response to UTI89 and 83972 is broadly conserved.  
9

10  
11  
12  
13 *Functional Annotation of DE Macrophage Genes*  
14

15 The consequences of gene expression changes associated with UPEC infection were  
16  
17 characterized by gene ontology (GO) and pathway (KEGG) enrichment analyses of DEG using the  
18  
19 DAVID program (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a). As shown in Supplementary Figure  
20  
21 2A, common highly enriched GO categories for upregulated genes included activation of inflammatory  
22  
23 responses; regulation of chemokine, cytokine, and interleukin-6 and -12 production; T-cell activation;  
24  
25 and regulation of transcription factor (TF) activity. Common biological processes associated with  
26  
27 downregulated genes included DNA-replication initiation, DNA packaging, nucleosome assembly and  
28  
29 organization, and cytokinesis. These pathways are consistent with the known activation of innate  
30  
31 immune responses by bacterial challenge (Rosenberger *et al.*, 2003, Mogensen, 2009, Portt *et al.*,  
32  
33 2011). As expected, the signaling pathways that were inferred, on the basis of gene expression  
34  
35 changes, to be regulated in macrophages upon UPEC infection showed common characteristics  
36  
37 between both UPEC strains. Signaling pathways associated with upregulated genes included  
38  
39 cytokine–cytokine receptor interaction, NOD-like receptor signaling, and Toll-like receptor (TLR)  
40  
41 signaling pathways (Figure 1D).  
42  
43  
44

45 We next independently validated both conservation and divergence in macrophage responses  
46  
47 to UTI89 versus 83972. Many of the well-validated TLR target genes such as cytokines (Il1a, Il1b, Il16)  
48  
49 and chemokines (Cxcl1, Ccl8) were similarly inducible by both UTI89 and 83972 in BMM (data not  
50  
51 shown). Although previous studies have demonstrated pathological roles for extracellular histones in  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 mouse models of sepsis (Xu *et al.*, 2009, Xu *et al.*, 2011), there is little known about the regulation of  
2 histone gene expression downstream of TLR signaling. This may reflect the fact that canonical histone  
3 mRNAs are not poly-adenylated, and so their regulated expression may not be captured by traditional  
4 microarray approaches. Interestingly, we found that a large suite of histone genes were  
5 downregulated in response to infection with either UPEC strain (Figure 2A), consistent with the  
6 inhibitory effect of TLR signaling on macrophage proliferation. This observation was validated for  
7 several individual histone genes using quantitative reverse transcription-polymerase chain reaction  
8 (RT-qPCR; Figure 2B). Our attempts to validate the small number of macrophage genes differentially  
9 regulated by UTI89 and 83972 (Figure 1B) using qPCR were less successful, however we did confirm  
10 differential regulation of the cystine/glutamate exchanger *Slc7a11*. Whereas *Slc7a11* mRNA  
11 expression was similarly upregulated by UTI89 and 83972 at 4 hpi in infected BMM, its expression  
12 remained elevated at 24 hpi in UTI89-infected BMMs but was significantly reduced at this time point in  
13 83972-infected BMMs (Figure 2C).

14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30 Clustering analysis separated DE macrophage genes into two major clusters (Figure 3A).  
31 Cluster 1 (Figure 3B), which contained 460 genes, was positively correlated with the profiles of the  
32 TFs *Arnt* (aryl hydrocarbon receptor nuclear translocator), *Myc* and *Pparg* (peroxisome proliferator-  
33 activated receptor gamma); and negatively correlated with the profiles of *Hif1a* (hypoxia-inducible  
34 factor 1-alpha) and *Stat3* (signal transducer and activator of transcription 3); and enriched for  
35 pathways associated with DNA replication, cell cycle, and systemic lupus erythematosus. Cluster 2  
36 (Figure 3B), which contained 217 genes, was positively correlated with the profiles of the TFs *Hif1a*,  
37 *Nfkb2* (nuclear factor of kappa light polypeptide gene enhancer in B cells 2), *Nr3c1* (nuclear receptor  
38 subfamily 3, group C, member 1) and *Stat3*; negatively correlated with the profile of *Myc*; and enriched  
39 for pathways associated with cytokine–cytokine receptor interaction, and NOD-like receptor, TLR,  
40 chemokine and Jak-STAT signaling. These results are in keeping with expectations; TLR signaling  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 activates *NF-κB* and *HIF* (Rossol *et al.*, 2011) and inactivates CSF-1 signaling. Various studies link  
3  
4 the biology of *CSF-1*, *Myc* and *Pparg* (Dey *et al.*, 2000, Bonfield *et al.*, 2008, Pello *et al.*, 2012), and as  
5  
6 expected, downregulated genes have an association with *Myc* and *Pparg*. That is, TLR signaling  
7  
8 switches off signaling via CSF-1, which itself can signal, in part, by *Myc*; hence, TLR signaling  
9  
10 downregulates *Myc* responses. Apart from the expected patterns, our analysis suggests a potential  
11  
12 association between inactivation of CSF-1 signaling and *Arnt* (partner for *Hif*) responses, which has  
13  
14 not been reported previously.  
15  
16  
17  
18  
19

20 *UPEC Transcriptional Responses upon Infection of BMMs*

21  
22 As was the case with the macrophage gene expression analysis, we initially investigated  
23  
24 transcriptional responses in both UPEC strains by performing dimensionality reduction analysis using  
25  
26 MCE (Cannistraci *et al.*, 2010, Cannistraci *et al.*, 2013). We identified specific differences in the  
27  
28 regulated expression of UPEC genes within the intramacrophage environment (Figure 4A), which  
29  
30 likely reflects the differential pathogenicity and capacity for intramacrophage survival of both strains  
31  
32 (Supplementary Figure 1) (Bokil *et al.*, 2011).  
33  
34

35 To gain insights into the global transcriptional changes that occur in UPEC during macrophage  
36  
37 infection, we applied the same method described above and performed pairwise comparisons  
38  
39 between controls (UPEC cultures alone) and treatments (co-cultures of UPEC-infected BMMs). Again,  
40  
41 we employed an FDR-adjusted *P*-value < 0.01 in at least one of the pairwise comparisons in our  
42  
43 dataset as the threshold for differential expression. In total, we identified 137 UTI89 genes and 33  
44  
45 83972 genes that were differentially regulated in BMMs. Surprisingly, an analysis of the  
46  
47 intramacrophage transcriptomic profile identified only 13 regulated genes that were common to both  
48  
49 strains, whereas 124 genes were uniquely regulated in UTI89, and 20 genes were uniquely regulated  
50  
51 in 83972 (Figure 4B). The UPEC transcriptional profiles also revealed a number of strain-specific  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 responses; a substantial number of UTI89 genes were upregulated at 2 and 4 hpi, whereas a much  
3  
4 smaller number of 83972 genes were upregulated at these time points (Figure 4C). This is consistent  
5  
6 with the initiation of a UTI89 transcriptional program that permits intramacrophage survival. In contrast  
7  
8 to the transcriptional responses observed for macrophage genes, few UPEC genes were  
9  
10 downregulated over the infection time course.  
11  
12  
13  
14

### 15 *Functional Annotation of DE UPEC Genes*

16  
17 The consequences of gene expression changes associated with UPEC infection were  
18  
19 characterized by GO and pathway (KEGG) enrichment analyses of DEG using the DAVID program  
20  
21 (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a). This again provided clear evidence of strain-specific  
22  
23 biological responses within the intramacrophage environment (Supplementary Figure 2B). Highly  
24  
25 enriched GO categories associated with the upregulated transcriptional program included genes  
26  
27 associated with chemotaxis and motility in UTI89 and genes involved in protein folding in 83972. We  
28  
29 did not identify any significantly enriched GO categories associated with downregulated genes in  
30  
31 common for the two strains. In keeping with the above findings, an analysis of the pathways activated  
32  
33 during UTI89 infection revealed enrichment for pathway terms associated with bacterial chemotaxis  
34  
35 and flagellar biosynthesis (Figure 4D). We did not observe significant enrichment for any pathway term  
36  
37 in the case of 83972.  
38  
39  
40  
41  
42

### 43 *UPEC Genes Associated with Intra-Macrophage Survival*

44  
45 We hypothesised that genes selectively upregulated by UTI89 may contribute to  
46  
47 intramacrophage survival. Such genes included those encoding flagella and those associated with  
48  
49 protection against oxidative stress. Notably, several genes encoding flagella-related proteins were  
50  
51 uniquely regulated in UTI89; these DE flagella genes included *flgA-F*, *flgK*, *flgL*, *fliC-E*, *fliQ*, *motA*, and  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 *motB*. Figure 5A shows the expression patterns of all flagella-related genes in UTI89 versus 83972,  
3  
4 revealing strong upregulation at 2 hours with a subsequent gradual decrease over time.  
5

6 A common mechanism used by bacterial pathogens to avoid host innate immune pathways is  
7  
8 to employ defense mechanisms against oxidative stress (Imlay, 2013). We therefore clustered the  
9  
10 expression patterns of all the OxyR regulon genes (Figure 5B) and the most strongly hydrogen  
11  
12 peroxide-induced genes in both UPEC strains (Figure 5C). Interestingly, the OxyR regulon, which  
13  
14 included alkyl hydroperoxide reductase subunit F (*ahpF*), *dps* (DNA starvation/stationary phase  
15  
16 protection protein), *grxA* (glutaredoxin 1), *trxC* (thioredoxin 2) and *yaaA*, was strongly upregulated in  
17  
18 UTI89, with the effect being apparent at 2 hpi and peaking at 4 hpi. Hydrogen peroxide-inducible  
19  
20 genes, which included *ahpF*, *dps*, *grxA*, heat-shock proteins/chaperones (*ibpA*, *ibpB*), *phoH*, *soxS*  
21  
22 (DNA-binding transcriptional dual regulator), *trxC* and *yaaA* showed a similar pattern, with genes being  
23  
24 significantly upregulated in UTI89 and less so in 83972. Differential regulation of *ahpF* in UTI89 versus  
25  
26 83972 within macrophages was validated by RT-qPCR (Figure 5D).  
27  
28  
29

30 Although UTI89 persists at 24 h within macrophages, only a relatively small percentage of  
31  
32 bacteria (< 5%) survive within BMM at 24 hpi compared to the bacterial loads at 2 hpi (Supplementary  
33  
34 Figure 1). We therefore reasoned that changes in UPEC gene expression at 24 hpi might be linked to  
35  
36 intramacrophage survival. We identified 22 genes that were highly upregulated (> 3-fold) by UTI89 at  
37  
38 24 hpi (Figure 6A). The most highly expressed of these included those encoding *ibpB* (encoding a  
39  
40 small heat shock protein); *pspACDE* (encoding the phage-shock protein system); *rpoE* and *rpoH*  
41  
42 (encoding sigma factors); *smpA* (encoding outer membrane lipoprotein); *yadR* (encoding the iron-  
43  
44 sulfur cluster insertion protein ErpA); *yceP* (encoding the biofilm formation regulatory protein BssS);  
45  
46 *yebG* (encoding DNA damage-inducible protein); and UTI89\_C2624 and UTI89\_C5162-3 (encoding  
47  
48 proteins of unknown function).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 Phage-shock-protein (Psp)-related genes, which are required for bacterial survival during  
3  
4 extracytoplasmic stress responses and changes in pH (Darwin, 2013), were significantly upregulated  
5  
6 in UTI89 compared to 83972 (Figure 6B). RT-qPCR confirmed the elevated expression of *pspA* and  
7  
8 *pspE* in UTI89 compared to 83972 at 24 h post-infection (Figure 6C). Finally, we validated the impact  
9  
10 of *pspA* for intramacrophage survival of UTI89 by constructing a UTI89*pspA* mutant (Figure 6D), and  
11  
12 testing it for intramacrophage survival during a 24 h infection time course in BMM. In this assay, the  
13  
14 UTI89*pspA* mutant was significantly reduced for intracellular survival compared to the wild type strain  
15  
16 (Figure 6E).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Discussion**

In this study, we determined the co-transcriptomic program of UPEC-infected primary mouse macrophages during an infection time course. The use of two UPEC strains that differed in their ability to survive in these cells enabled the identification of both common and UPEC strain-specific responses. To our knowledge, this is one of the first RNA-Seq studies that have simultaneously measured the transcriptomes of both the host and pathogen during an infection (Humphrys *et al.*, 2013). Using only open-source tools, we developed a computational framework that was capable of successfully separating, annotating, and quantifying the mammalian and bacterial transcriptomes. Whereas previous studies (Mysorekar *et al.*, 2002, Bower *et al.*, 2009, Hagan *et al.*, 2010, Duell *et al.*, 2012) have been limited to microarrays, the RNA-Seq analysis reported here provides a more sensitive, comprehensive, and unbiased coverage of the entire transcriptome. We achieved an average sequencing depth of approximately 24 million tags per library and identified 677 BMM and 157 UPEC genes that were differentially expressed following UPEC infection. By mapping our RNA-Seq tag data onto transcript databases and genomic sequences, we were able to identify genes that were regulated upon UPEC challenge.

MCE confirmed the distinct temporal cascade of BMM responses to UPEC as well as differences in transcriptional programs of the two UPEC strains. The sets of DEG and regulatory processes and pathways identified in both BMMs and UPEC strains suggest coordinated expression and mutual influences. Our study detected 603 DEG in BMMs that were common to infection with UTI89 and 83972, accounting for the vast majority of changes in BMM gene expression. This was not unexpected, given that both UTI89 and 83972 present pathogen-associated molecular patterns, such as lipopolysaccharide that is recognized by TLR4. Macrophage-expressed genes that showed differences in regulation in response to these two strains were a very minor component of the total



signature and their significance remains to be determined. Despite the general conservation of the BMM response against UPEC infection, we nonetheless expected to detect some selective BMM responses at 24 hpi for UTI89 compared with 83972, since the former condition reflects macrophages that are continuing to cope with intracellular UPEC, whereas the latter corresponds to macrophages that have cleared the infection. Consistent with this supposition, we found that at 24 hpi, BMM responses showed a greater divergence between UTI89 and 83972 (47.8% overlap) compared to earlier time points where this difference was not as evident. Of interest in this regard, was our validation of differential regulation of the cationic amino acid cysteine/glutamate antiporter Slc7a11, which is required for glutamate uptake and glutathione synthesis (Bannai, 1986, Hayes *et al.*, 1999, Pompella *et al.*, 2003, Shih *et al.*, 2006). This gene showed a sustained upregulation at 24 hpi following infection with UTI89, whereas it was only transiently upregulated after infection with 83972. Given that macrophage antimicrobial responses typically involve subjecting pathogens to oxidative stress, the sustained expression of Slc7a11 upon infection with UTI89 may be required to maintain glutathione levels and cytoprotection during stress responses.

Our data is consistent with previous reports showing inducible cytokine and chemokine expression, as well as activation of pro-survival pathways (Sester *et al.*, 1999, Sester *et al.*, 2006), during bacterial infection and/or TLR stimulation of macrophages. Moreover, the downregulation of DNA replication and cell cycle genes (Figure 1D) is consistent with the known inhibitory effects of TLR4 signalling on proliferation of cycling macrophages (Sester *et al.*, 1999, Sester *et al.*, 2006). We also found that the genes encoding histones H1, H2, and H4 were dramatically downregulated at 24 hpi, and we further confirmed the regulated expression of several of these using RT-qPCR. A member of the H2A histone family, *Hist2h2aa1*, was also significantly downregulated at 24 hpi *in vivo* in mouse bladder colonized with UPEC CFT073 (Tan *et al.*, 2012), which is consistent with these findings. Interestingly, previous studies have reported a pathological role for extracellular histones during LPS-

1 induced septic shock (Xu *et al.*, 2009, Li *et al.*, 2011). Xu *et al.* (2011) revealed that antibodies against  
2 extracellular histones rescued animals from LPS-mediated death (Xu *et al.*, 2011), and a previous  
3 study showed that extracellular histones mediate endothelial dysfunction, organ failure, and death  
4 during sepsis (Semeraro *et al.*, 2011). Our findings could thus either reflect a host attempt to reduce  
5 inflammatory responses upon cell death and histone release, or the consequence of growth-inhibitory  
6 effects of TLR agonists on proliferating macrophages, which would also be expected to lead to  
7 downregulated histone expression.  
8  
9

10 A transcription factor binding site (TFBS) analysis of the promoter sequences of the two BMM  
11 DE gene clusters further showed significant enrichment of seven motifs associated with TFs that have  
12 key roles in macrophage functions. The cluster containing inducible inflammation-related genes  
13 correlated positively with the expression of *Hif1a*, a key pro-inflammatory transcription factor that  
14 drives macrophage inflammatory responses and is upregulated in UPEC-infected mouse bladder at 2  
15 hpi and 24 hpi (Duell *et al.*, 2012, Tan *et al.*, 2012); *Nfkb2*, which is activated by a wide variety of  
16 stimuli such as cytokines, oxidant-free radicals and bacterial or viral products; the glucocorticoid  
17 receptor *Nr3c1*, which up-regulates the expression of anti-inflammatory genes and/or represses the  
18 expression of pro-inflammatory genes; and *Stat3*, a transcriptional activator stimulated in response to  
19 cytokines and growth factors. *In vivo*, both *Nfkb2* and *Stat3* are immediately upregulated in the bladder  
20 following UPEC infection (Duell *et al.*, 2012). On the other hand, the cluster containing downregulated  
21 cell cycle-related genes correlated positively with the expression of *Arnt*, which is involved in the  
22 induction of several enzymes that participate in xenobiotic metabolism; *Myc*, which is required for cell  
23 proliferation in response to mitogenic stimuli such as CSF-1; and *Pparg*, which regulates fatty acid  
24 storage and glucose metabolism. *Myc*, *Pparg*, and *Csf1* are all differentially expressed in bladders of  
25 mice infected with UPEC (Duell *et al.*, 2012) supporting the *in vivo* relevance of these findings.  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 In contrast to the general conserved pattern of BMM responses to both UPEC strains, the  
3 individual transcriptional responses of the two strains in BMMs differed markedly. This is consistent  
4 with their contrasting survival patterns in macrophages. MCE confirmed the different transcriptional  
5 programs of the two UPEC strains, revealing 124 UTI89-specific DEG, 20 83972-specific DEG, and 13  
6 DEG that were commonly regulated in both pathogens upon infection of BMMs. Given the different  
7 survival patterns of both UPEC strains in macrophages, we expected to detect increased expression  
8 of genes specific to UTI89 within macrophages at 24 hpi.  
9  
10  
11  
12  
13  
14  
15  
16

17 Previous studies have shown that flagella contribute to the virulence of a number of pathogenic  
18 species (Tomich *et al.*, 2002). Some bacteria, for example *Salmonella enterica* Typhimurium and  
19 *Yersinia enterocolitica*, use flagella for invasion of epithelial cells (McNally *et al.*, 2007, Ibarra *et al.*,  
20 2010). UPEC also employ flagella for the invasion of renal collecting duct cells (Pichon *et al.*, 2009),  
21 but flagella-enhanced uptake of UPEC by macrophages has not been investigated. Our co-  
22 transcriptomic analysis revealed the upregulation of multiple genes associated with flagella  
23 biosynthesis. This could reflect inducible gene expression in the intramacrophage environment or  
24 selective uptake by macrophages of a sub-population of flagella-expressing UPEC, which would  
25 generate a similar gene expression profile. The acute upregulation at 2 hpi and subsequent  
26 downregulation at 24 hpi would suggest the latter may be the case.  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

39 Murine macrophages employ both rapidly produced reactive oxygen species (ROS) and  
40 reactive nitrogen species (RNS), produced in a delayed fashion, as bacterial clearance strategies  
41 (Flannagan *et al.*, 2009). The observation that UPEC strain UTI89 survives in macrophages suggests  
42 that it can overcome these pathways. Although the genes that respond to ROS have been well  
43 characterized in *E. coli* K-12, their expression during UPEC infection and their role in intramacrophage  
44 survival have not yet been studied. Our transcriptomic approach identified UPEC genes that are likely  
45 involved in defense against ROS and RNS. Disturbances in the normal redox state of cells can cause  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 toxic effects through the production of peroxides and free radicals that damage all components of the  
3  
4 cell, including proteins, lipids, and DNA. An earlier study confirmed that the peroxide response  
5  
6 regulator OxyR activates most of the genes that are highly induced by hydrogen peroxide (Zheng *et*  
7  
8 *al.*, 2001). Our results show that the response of UTI89 to oxidative stress (upregulated expression of  
9  
10 *dps*, *grxA*, *ahpF*, *soxS*, *trxC*, *ibpA*, and *ibpB*) is more robust than that of 83972. These differences  
11  
12 imply that part of the UTI89 intramacrophage survival strategy includes robust protection against  
13  
14 oxidative stress.  
15  
16

17  
18 Several UTI89 genes were upregulated in BMM at 24 hpi, and we selected genes encoding the  
19  
20 Psp system for further functional analysis. The Psp system responds to extracytoplasmic stress and  
21  
22 contributes to the virulence of several pathogens, including *Y. enterocolitica* and *S. Typhimurium*  
23  
24 (Karlinsey *et al.*, 2010, Yamaguchi *et al.*, 2012). PspA, the master effector of the Psp system,  
25  
26 mediates its response via a dual mechanism: (i) binding to the transcriptional regulator PspF and  
27  
28 preventing it from activating the transcription of *pspACDE* in the absence of extracytoplasmic stress,  
29  
30 and (ii) binding to the cytoplasmic membrane-localized proteins PspB and PspC in the presence of  
31  
32 extracytoplasmic stress, thus releasing PspF to induce *pspACDE* transcription (Yamaguchi *et al.*,  
33  
34 2013). We confirmed a role for the Psp system in UPEC by constructing a *pspA* mutant (UTI89*pspA*)  
35  
36 and demonstrating reduced intramacrophage survival of this mutant. Despite this statistically  
37  
38 significant reduction in survival within macrophages, the biological significance of this effect is at this  
39  
40 stage unknown. Inactivation of other stress response systems on the *pspA* mutant background may be  
41  
42 required to reveal a more striking phenotype.  
43  
44

45  
46 In summary, we have demonstrated the capacity to employ co-transcriptomics to study host-  
47  
48 pathogen interactions. Our novel approach revealed new insights into the mechanisms used by UPEC  
49  
50 to avoid macrophage responses and persist in the intra-macrophage environment, and has identified  
51  
52 multiple target genes for further functional studies. Finally, this manuscript is one of the first to  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 successfully evaluate the expression profiles of two organisms from the same sample using RNA  
3  
4 sequencing, and will make an excellent resource for other studies that aim to perform similar analyses  
5  
6 in different host-pathogen systems.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2 **Experimental Procedures**

3  
4  
5  
6 *TIER I: In vitro Infection Assays*

7  
8 Ethics Statement and Animal Experimentation

9  
10 A University of Queensland institutional animal ethics committee approved all animal  
11 experimentations. Female C57BL/6 mice (6–8 weeks old) were purchased from the Animal Resources  
12 Center, Australia. Murine bone marrow-derived macrophages (BMM) were generated by the *in vitro*  
13 differentiation of bone marrow cells from C57BL/6 mice on bacteriological plastic plates in the  
14 presence of 10,000 U ml<sup>-1</sup> recombinant human CSF-1 (a gift from Chiron) for 6 days, after which cells  
15 were harvested and replated onto tissue culture plastic in the presence of CSF-1 for infection on day  
16 7. BMM were maintained at 37°C (5% CO<sub>2</sub>) in RPMI 1640 supplemented with 2mM L-glutamine  
17 (GlutaMAX), 10% heat-inactivated fetal bovine serum and 50 U ml<sup>-1</sup> penicillin and 50 µg ml<sup>-1</sup>  
18 streptomycin (Life Technologies).  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Culture of Bacterial Strains and Infection of Mouse BMMs

33  
34 *E. coli* UTI89 is a well-characterized cystitis isolate (Mulvey *et al.*, 2001). *E. coli* 83972, which  
35 was carried without symptoms by a young female, was originally isolated from the urine of this  
36 individual (Andersson *et al.*, 1991, Roos *et al.*, 2006). For macrophage infection assays, UPEC strains  
37 were cultured statically in Luria-Bertani (LB) broth at 37°C overnight. Type 1 fimbriae expression was  
38 assessed by yeast cell agglutination prior to infection as previously described (Schembri *et al.*, 2000).  
39 Bacterial cells were centrifuged and washed in phosphate buffered saline (PBS), and then  
40 resuspended in antibiotic-free media at a concentration of 2 x 10<sup>8</sup> CFU ml<sup>-1</sup>. Viable CFU counts of  
41 bacterial inocula were routinely confirmed in every infection assay by serial dilution and plating on LB  
42 agar.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Intramacrophage survival assays were performed essentially as described previously (Bokil *et al.*, 2011). Briefly, following overnight adherence in antibiotic-free media, BMMs were infected for 1h with bacteria at an MOI of 10. Extracellular bacteria were killed by washing twice in 200  $\mu\text{g ml}^{-1}$  gentamicin, followed by 1h incubation in media containing the same gentamicin concentration. Subsequent exclusion of extracellular bacteria for the duration of the experiment was performed by incubation in 20  $\mu\text{g ml}^{-1}$  gentamicin. At appropriate time points (1h, 2h and 24h), cells were washed twice with antibiotic-free media, and then lysed with PBS/0.01% Triton X-100. Lysates were cultured on LB agar plates overnight at 37°C and colony counts were used to assess intra-cellular bacterial loads. For standard infection assays, complete exclusion of viable extracellular bacteria was confirmed by performing colony counts on culture supernatants.

## *TIER II: RNA Preparation and Sequencing*

### Macrophage Infection and mRNA Isolation, Enrichment, and Purification

Control macrophages, control bacteria and infected macrophages were incubated for an additional 1, 3 and 23 h at 37°C in a humidified 5% CO<sub>2</sub> atmosphere, after the initial 1 h infection. Cells were then washed twice with antibiotic-free media, and then lysed on ice for RNA isolation and purification (RNeasy; Qiagen, Germantown, MD, USA). Microbial total RNA in co-culture samples was enriched (MICROBEnrich, Ambion). rRNA was removed from all purified RNA samples using kits targeting mammalian and Gram-negative bacterial rRNAs (Ribo-Zero; Epicenter, Madison, WI, USA). Prior to sequencing, all samples were further quantified and examined for protein and reagent contamination using a Nanodrop ND-1000 spectrophotometer. RNA samples for analysis were selected based on a 28S/18S rRNA band intensity of 2:1, a spectroscopic  $A_{260}/A_{280}$  nm ratio of 1.8–2.0, and an  $A_{260}/A_{230}$  nm ratio > 1.5.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

RNA Sequencing

Next-generation sequencing analyses were performed for two biological replicates on an Illumina Cluster Station and the Illumina HiSeq 2000 System using primarily reagents from the Illumina Gene Expression Sample Preparation Kit and the Illumina Sequencing Chip (Flowcell; Illumina, San Diego, CA, USA). Sequence tags were prepared using the Digital Gene Expression Tag Profiling Kit (Illumina), according to the Illumina protocol.

*TIER III: Data Pre-processing*

Image analysis, base calling, and quality calibration were performed using the Solexa Automated Pipeline. Quality control of RNA-Seq reads were preprocessed by a custom java script. All sequences generated have a length of 101 bases and have been submitted to the BioProject database of NCBI under BioProject ID: PRJNA256028.

*TIER IV: Alignment and Differential Gene Expression Analysis*

Alignment of Reads

Bowtie indexes were created for the mouse (version 37.1) and the two *E. coli* strains (UTI89 and 83972) using the bowtie-build algorithm and reference sequences from the GenBank database. Our protocol began with mapping of the raw RNA-Seq reads (fastq files) to the reference genomes using TopHat. TopHat uses Bowtie as an alignment engine and breaks up reads that Bowtie cannot align on its own into smaller pieces (Kim *et al.*, 2011). Using the standard parameters, we mapped both reads of our paired-end libraries. All simulations were performed using a 30-core, high-memory node cluster system; total computation duration was 3 hours.

Transcript Annotation



After running TopHat, the resulting alignment files were provided to Cufflinks to generate a transcriptome assembly for each condition. During this analysis step, adapter tags; mitochondrial sequences; poly A, poly C, and phiX sequences; and remaining ribosomal sequences were filtered out. Estimated normalized expression levels were reported in Fragments (i.e., reads) Per Kilobase of exon per Million mapped reads (FPKM). These assemblies were compared with annotation files using the Cuffcompare utility, which is included in the Cufflinks package (Roberts *et al.*, 2011, Trapnell *et al.*, 2012).

### Differential Expression Analysis

The reads and assemblies were imported to Cuffdiff, which calculates expression levels and tests the statistical significance of observed changes. For comparison of DEG across samples, the number of raw clean tags in each library was normalized to FPKM using the Cufflinks package. The minimum number of alignments in a locus needed to test for significance of changes in that locus between samples was set to 50 fragment alignments. If no testing was performed, changes in the locus were deemed insignificant, and the changes observed in the locus did not contribute to corrections for multiple testing. The Cuffdiff output files were then imported to cummeRbund, which plots abundance and differential expression results as commonly used expression plots for quality control (Trapnell *et al.*, 2012).

### *TIER V: Functional Analysis*

#### Dimensionality Reduction

Dimensionality reduction is necessary for exploring the relationships between conditions in our experiment. The MCE method performs a nonlinear dimension reduction by embedding high-dimensional data points into a lower-dimensional space using the minimum curvilinear kernel in

1 combination with multidimensional scaling (MDS) (Cannistraci *et al.*, 2010) or in alternative the  
2  
3 singular value decomposition (Cannistraci *et al.*, 2013). The nonlinear data distances for MDS or SVD  
4  
5 were computed and stored in the minimum curvilinear kernel as the traversal distances over the  
6  
7 minimum spanning tree between the data points (in our study the samples' conditions) in the  
8  
9 multidimensional space (in our case the gene space). The minimum spanning tree was constructed  
10  
11 from the Pearson correlation-based distances between the samples (Cannistraci *et al.*, 2010):  
12  
13

14  
15 
$$\text{correlation\_based\_distance}(x,y) = 1 - \text{Pearson\_correlation}(\text{Sample}_x, \text{Sample}_y)$$
  
16

17 MCE is a parameter-free projection algorithm that was shown to be particularly effective in  
18  
19 discriminating classes in small-*n* (samples: here conditions), large-*m* (features: here gene  
20  
21 expressions) datasets using only the first dimension of embedding (Cannistraci *et al.*, 2010). The fact  
22  
23 that our datasets have  $n \ll m$  led us to adopt MCE algorithm for unsupervised analysis of the patterns  
24  
25 present between the different sample conditions.  
26  
27

28  
29  
30 Clustering of Genes  
31

32 Genes with similar expression patterns often serve overlapping functions. Accordingly, the  
33  
34 optimal number of clusters in the dataset was determined by performing a cluster analysis of gene  
35  
36 expression patterns using the R package NbClust. Selected lists of expression profiles of DE genes  
37  
38 were compiled for each hypothesis tested and clustered using the Ward's methodology.  
39  
40  
41

42  
43 GO and Pathway-enrichment analyses of DEG  
44

45 Genes involved in common biological processes or pathways tend to show overlapping  
46  
47 expression profiles. In gene expression profiling analyses, significantly enriched GO terms and  
48  
49 pathways were identified by mapping all DEG to terms in the GO and KEGG databases by applying  
50  
51 two-sided Fisher's exact and  $\chi^2$  tests, respectively (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a).  
52  
53  
54  
55  
56  
57  
58  
59  
60

P-values were corrected by calculating the FDR, and only GO and pathway terms with a FDR < 0.01 were chosen.

### Identification of key BMM TFs

Promoter sequences of all DEG were retrieved using the RSAT and were further input into the RSAT matrix-scan tool along with mouse-related JASPAR matrices for TFBS prediction (Thomas-Chollier *et al.*, Thomas-Chollier *et al.*, 2008, Turatsinze *et al.*, 2008). The RSAT output was filtered using an adjusted P-value < 0.05 as a cut-off, and lists of the most significant TFBSs and their known corresponding TFs were compiled. The expression profiles of the mouse DEG were clustered and each cluster was correlated with TFs profiles using Pearson correlation in R. Finally, the clusters were annotated using GO to fully elucidate the molecular processes in which each TF was involved.

### Screening for strain-specific UPEC gene expression patterns

Flagella-related gene lists were compiled based on Macnab's review on bacterial flagellar assembly (Macnab, 2003). A list of 35 genes related to the flagellar apparatus was used as background for screening our UPEC datasets. For OxyR regulon and hydrogen peroxide-induced genes, gene lists were similarly compiled using previously published data on the response of bacteria to hydrogen peroxide (Zheng *et al.*, 2001). Finally, lists of genes involved in Psp regulation were compiled, and our datasets were screened for their expression patterns. The expression patterns of the screened genes were further clustered as described above, and all results were visualized using pheatmap package in R.

### Identification of UPEC Genes Associated with Intramacrophage Survival

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Putative bacterial genes associated with intramacrophage survival were considered as those that remained highly upregulated at 24 hpi. Bacterial genes upregulated at this time point were compiled and filtered based on their significance of expression. All DEG were filtered using an FDR-adjusted P-value of the test statistic  $< 0.01$  and a  $\log_2$  fold change  $> 3$ . The filtered lists of DEG enabled us to screen for survival genes, cluster their expression, and further annotate them using the GO database to better understand the biological processes they regulated.

*TIER VI: Gene Validation*

cDNA synthesis, primer design and RT-qPCR

cDNA was synthesized using a SuperScript III First-Strand Synthesis kit (Invitrogen). qPCR was then performed on a 7900HT Fast Real-Time PCR system (Applied Biosystems) using RNA samples from two independent biological replicates, similar to those employed for the RNA-Seq experiments. TaqMan Fast Universal PCR Master Mix 2X (Applied Biosystems) was used for BMM gene validation, and SYBR Green Master Mix (Applied Biosystems) was used for UPEC gene validation. Each cDNA was analyzed in triplicate, after which the average threshold cycle (Ct) per sample was calculated. Raw data were processed with qBase Plus software (Biogazelle), which performs downstream processing of qPCR data. The geNorm algorithm, integrated in the qBase Plus package, was used for determining the optimal number and identity of reference genes needed to normalize the data in both BMM (Actb and Polr2a) and UPEC (gapA and purC) qPCR libraries. Relative expression levels were calculated with the  $2^{-\Delta\Delta Ct}$  method ( $\Delta Ct$  is the difference in Ct between the reference genes and the target gene products); the average Ct value for all genes was used to correct for differences in cDNA input. Other statistical procedures were performed with the R program. All steps, from the experimental design to bioinformatic analysis and gene validation, are summarized in Supplementary Figure 3.

### Construction of UTI89 $\psi$ spA deletion mutant

Chromosomal DNA purification, PCR and DNA sequencing of PCR products was performed as previously described (Allsopp *et al.*, 2010). The  $\psi$ spA gene was mutated in UTI89 using the  $\lambda$ -Red mediated homologous recombination method with some modifications (Datsenko *et al.*, 2000, Allsopp *et al.*, 2012). Briefly, a three-step PCR procedure was employed to generate a DNA fragment comprising the chloramphenicol cassette from plasmid pKD3 and two 500-bp fragments homologous to the flanking regions of the  $\psi$ spA gene. The following primers were used: 5376\_UTI89 $\psi$ spA FwUP (5'-gccgtagcgagttcatca) and 5377\_UTI89 $\psi$ spA RvUP (5'-ggaataggaactaaggaggaagcgttgatgttggcatt), 5378\_UTI89 $\psi$ spA Fwdn (5'-cctacacaatcgctcaagacgccgaactgaaagccgat) and 5379\_UTI89 $\psi$ spA Rvdn (5'-taaacagcgccagaccga) to generate the 500bp homology arms; 3746-Cm.3a (5'-tcctccttagttcctattcc) and 3747-Cm.4a (5'-gtcttgagcgattgtgtagg) to generate the chloramphenicol resistance gene fragment. This DNA fusion product was electroporated into UTI89 harboring plasmid pKD46, and chloramphenicol resistant mutants were selected and confirmed by PCR (using primers 5375\_UTI89 $\psi$ spA FwSc: 5'-tcgtcgcgcataccaacc and 5380\_UTI89 $\psi$ spA Rvsc: 5'-acttcatccagcaattcgc). The UTI89 $\psi$ spA mutant was confirmed by sequencing.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Acknowledgments**

This work was supported by grants from the National Health and Medical Research Council (NHMRC) of Australia (APP1005315 and APP1068593). MJS is supported by an Australian Research Council (ARC) Future Fellowship (FT100100657), as well as an honorary NHMRC Senior Research Fellowship (APP1003470). MAS and GCU are supported by ARC Future Fellowships (FT100100662 and FT110101048, respectively), MT is supported by an ARC Discovery Early Career Researcher Award (DE130101169), and SAB is supported by an NHMRC Career Development Fellowship (APP1090456).

## References

- Allsopp, L.P., Beloin, C., Ulett, G.C., Valle, J., Totsika, M., Sherlock, O., *et al.* (2012). Molecular characterization of UpaB and UpaC, two new autotransporter proteins of uropathogenic *Escherichia coli* CFT073. *Infection and immunity* **80**, 321-332.
- Allsopp, L.P., Totsika, M., Tree, J.J., Ulett, G.C., Mabbett, A.N., Wells, T.J., *et al.* (2010). UpaH is a newly identified autotransporter protein that contributes to biofilm formation and bladder colonization by uropathogenic *Escherichia coli* CFT073. *Infection and immunity* **78**, 1659-1669.
- Andersson, P., Engberg, I., Lidin-Janson, G., Lincoln, K., Hull, R., Hull, S. and Svanborg, C. (1991). Persistence of *Escherichia coli* bacteriuria is not determined by bacterial adherence. *Infection and immunity* **59**, 2915-2921.
- Bannai, S. (1986). Exchange of cystine and glutamate across plasma membrane of human fibroblasts. *J Biol Chem* **261**, 2256-2263.
- Bokil, N.J., Totsika, M., Carey, A.J., Stacey, K.J., Hancock, V., Saunders, B.M., *et al.* (2011). Intramacrophage survival of uropathogenic *Escherichia coli*: differences between diverse clinical isolates and between mouse and human macrophages. *Immunobiology* **216**, 1164-1171.
- Bonfield, T.L., Thomassen, M.J., Farver, C.F., Abraham, S., Koloze, M.T., Zhang, X., *et al.* (2008). Peroxisome proliferator-activated receptor-gamma regulates the expression of alveolar macrophage colony-stimulating factor. *Journal of immunology* **181**, 235-242.
- Bower, J.M., Gordon-Raagas, H.B. and Mulvey, M.A. (2009). Conditioning of uropathogenic *Escherichia coli* for enhanced colonization of host. *Infect Immun* **77**, 2104-2112.
- Cannistraci, C.V., Alanis-Lobato, G. and Ravasi, T. (2013). Minimum curvilinearity to enhance topological prediction of protein interactions by network embedding. *Bioinformatics* **29**, i199-209.
- Cannistraci, C.V., Ravasi, T., Montevecchi, F.M., Ideker, T. and Alessio, M. (2010). Nonlinear dimension reduction and clustering by Minimum Curvilinearity unfold neuropathic pain and tissue embryological classes. *Bioinformatics* **26**, i531-539.
- Darwin, A.J. (2013). Stress relief during host infection: The phage shock protein response supports bacterial virulence in various ways. *PLoS pathogens* **9**, e1003388.
- Datsenko, K.A. and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 6640-6645.
- Dey, A., She, H., Kim, L., Boruch, A., Guris, D.L., Carlberg, K., *et al.* (2000). Colony-stimulating factor-1 receptor utilizes multiple signaling pathways to induce cyclin D2 expression. *Molecular biology of the cell* **11**, 3835-3848.
- Duell, B.L., Carey, A.J., Tan, C.K., Cui, X., Webb, R.I., Totsika, M., *et al.* (2012). Innate transcriptional networks activated in bladder in response to uropathogenic *Escherichia coli* drive diverse biological pathways and rapid synthesis of IL-10 for defense against bacterial urinary tract infection. *Journal of immunology* **188**, 781-792.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Eden, C.S., Hanson, L.A., Jodal, U., Lindberg, U. and Akerlund, A.S. (1976). Variable adherence to normal human urinary-tract epithelial cells of *Escherichia coli* strains associated with various forms of urinary-tract infection. *Lancet* **1**, 490-492.

Flannagan, R.S., Cosio, G. and Grinstein, S. (2009). Antimicrobial mechanisms of phagocytes and bacterial evasion strategies. *Nature reviews. Microbiology* **7**, 355-366.

Foxman, B. (2010). The epidemiology of urinary tract infection. *Nat Rev Urol* **7**, 653-660.

Funfstuck, R., Tschape, H., Stein, G., Kunath, H., Bergner, M. and Wessel, G. (1986). Virulence properties of *Escherichia coli* strains in patients with chronic pyelonephritis. *Infection* **14**, 145-150.

Hacker, J. and Kaper, J.B. (2000). Pathogenicity islands and the evolution of microbes. *Annu Rev Microbiol* **54**, 641-679.

Hagan, E.C., Lloyd, A.L., Rasko, D.A., Faerber, G.J. and Mobley, H.L. (2010). *Escherichia coli* global gene expression in urine from women with urinary tract infection. *PLoS Pathog* **6**, e1001187.

Hannan, T.J., Totsika, M., Mansfield, K.J., Moore, K.H., Schembri, M.A. and Hultgren, S.J. (2012). Host-pathogen checkpoints and population bottlenecks in persistent and intracellular uropathogenic *Escherichia coli* bladder infection. *FEMS Microbiol Rev* **36**, 616-648.

Hayes, J.D. and McLellan, L.I. (1999). Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defence against oxidative stress. *Free radical research* **31**, 273-300.

Hegedus, Z., Zakrzewska, A., Agoston, V.C., Ordas, A., Racz, P., Mink, M., *et al.* (2009). Deep sequencing of the zebrafish transcriptome response to mycobacterium infection. *Mol Immunol* **46**, 2918-2930.

Horvath, D.J., Jr., Dabdoub, S.M., Li, B., Vanderbrink, B.A. and Justice, S.S. (2012). New paradigms of urinary tract infections: Implications for patient management. *Indian J Urol* **28**, 154-158.

Huang da, W., Sherman, B.T. and Lempicki, R.A. (2009a). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13.

Huang da, W., Sherman, B.T. and Lempicki, R.A. (2009b). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57.

Huang, Q., Dong, S., Fang, C., Wu, X., Ye, T. and Lin, Y. (2012). Deep sequencing-based transcriptome profiling analysis of *Oryzias melastigma* exposed to PFOS. *Aquat Toxicol* **120-121**, 54-58.

Humphrys, M.S., Creasy, T., Sun, Y., Shetty, A.C., Chibucos, M.C., Drabek, E.F., *et al.* (2013). Simultaneous transcriptional profiling of bacteria and their host cells. *PLoS One* **8**, e80597.

Hunstad, D.A. and Justice, S.S. (2010). Intracellular lifestyles and immune evasion strategies of uropathogenic *Escherichia coli*. *Annu Rev Microbiol* **64**, 203-221.



- Ibarra, J.A., Knodler, L.A., Sturdevant, D.E., Virtaneva, K., Carmody, A.B., Fischer, E.R., *et al.* (2010). Induction of Salmonella pathogenicity island 1 under different growth conditions can affect Salmonella-host cell interactions in vitro. *Microbiology* **156**, 1120-1133.
- Imlay, J.A. (2013). The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nature reviews. Microbiology* **11**, 443-454.
- Ingersoll, M.A., Kline, K.A., Nielsen, H.V. and Hultgren, S.J. (2008). G-CSF induction early in uropathogenic Escherichia coli infection of the urinary tract modulates host immunity. *Cellular microbiology* **10**, 2568-2578.
- Jager, D., Sharma, C.M., Thomsen, J., Ehlers, C., Vogel, J. and Schmitz, R.A. (2009). Deep sequencing analysis of the Methanosarcina mazei Go1 transcriptome in response to nitrogen availability. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 21878-21882.
- Johnson, J.R. (1991). Virulence factors in Escherichia coli urinary tract infection. *Clin Microbiol Rev* **4**, 80-128.
- Kaijser, B. and Ahlstedt, S. (1977). Protective capacity of antibodies against Escherichia coli and K antigens. *Infection and immunity* **17**, 286-289.
- Karlinsey, J.E., Maguire, M.E., Becker, L.A., Crouch, M.L. and Fang, F.C. (2010). The phage shock protein PspA facilitates divalent metal transport and is required for virulence of Salmonella enterica sv. Typhimurium. *Molecular microbiology* **78**, 669-685.
- Kim, D. and Salzberg, S.L. (2011). TopHat-Fusion: an algorithm for discovery of novel fusion transcripts. *Genome Biol* **12**, R72.
- Klemm, P., Hancock, V. and Schembri, M.A. (2007). Mellowing out: adaptation to commensalism by Escherichia coli asymptomatic bacteriuria strain 83972. *Infect Immun* **75**, 3688-3695.
- Lau, M.E., Loughman, J.A. and Hunstad, D.A. (2012). YbcL of uropathogenic Escherichia coli suppresses transepithelial neutrophil migration. *Infection and immunity* **80**, 4123-4132.
- Li, Y., Liu, B., Fukudome, E.Y., Lu, J., Chong, W., Jin, G., *et al.* (2011). Identification of citrullinated histone H3 as a potential serum protein biomarker in a lethal model of lipopolysaccharide-induced shock. *Surgery* **150**, 442-451.
- Lindberg, U. (1975). Asymptomatic bacteriuria in school girls. V. The clinical course and response to treatment. *Acta paediatrica Scandinavica* **64**, 718-724.
- Lindberg, U., Bjure, J., Haugstvedt, S. and Jodal, U. (1975a). Asymptomatic bacteriuria in schoolgirls. III. Relation between residual urine volume and recurrence. *Acta paediatrica Scandinavica* **64**, 437-440.
- Lindberg, U., Claesson, I., Hanson, L.A. and Jodal, U. (1975b). Asymptomatic bacteriuria in schoolgirls. I. Clinical and laboratory findings. *Acta paediatrica Scandinavica* **64**, 425-431.
- Lindberg, U., Hanson, L.A., Jodal, U., Lidin-Janson, G., Lincoln, K. and Olling, S. (1975c). Asymptomatic bacteriuria in schoolgirls. II. Differences in escherichia coli causing asymptomatic bacteriuria. *Acta paediatrica Scandinavica* **64**, 432-436.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Llorens, F., Hummel, M., Pastor, X., Ferrer, A., Pluvinet, R., Vivancos, A., *et al.* (2011). Multiple platform assessment of the EGF dependent transcriptome by microarray and deep tag sequencing analysis. *BMC Genomics* **12**, 326.

Mabbett, A.N., Ulett, G.C., Watts, R.E., Tree, J.J., Totsika, M., Ong, C.L., *et al.* (2009). Virulence properties of asymptomatic bacteriuria Escherichia coli. *International journal of medical microbiology : IJMM* **299**, 53-63.

Macnab, R.M. (2003). How bacteria assemble flagella. *Annu Rev Microbiol* **57**, 77-100.

McNally, A., La Ragione, R.M., Best, A., Manning, G. and Newell, D.G. (2007). An aflagellate mutant Yersinia enterocolitica biotype 1A strain displays altered invasion of epithelial cells, persistence in macrophages, and cytokine secretion profiles in vitro. *Microbiology* **153**, 1339-1349.

Mogensen, T.H. (2009). Pathogen recognition and inflammatory signaling in innate immune defenses. *Clinical microbiology reviews* **22**, 240-273, Table of Contents.

Morozova, O. and Marra, M.A. (2008). Applications of next-generation sequencing technologies in functional genomics. *Genomics* **92**, 255-264.

Mulvey, M.A., Schilling, J.D. and Hultgren, S.J. (2001). Establishment of a persistent Escherichia coli reservoir during the acute phase of a bladder infection. *Infection and immunity* **69**, 4572-4579.

Mysorekar, I.U. and Hultgren, S.J. (2006). Mechanisms of uropathogenic Escherichia coli persistence and eradication from the urinary tract. *Proc Natl Acad Sci U S A* **103**, 14170-14175.

Mysorekar, I.U., Mulvey, M.A., Hultgren, S.J. and Gordon, J.I. (2002). Molecular regulation of urothelial renewal and host defenses during infection with uropathogenic Escherichia coli. *J Biol Chem* **277**, 7412-7419.

Nie, Q., Sandford, E.E., Zhang, X., Nolan, L.K. and Lamont, S.J. (2012). Deep sequencing-based transcriptome analysis of chicken spleen in response to avian pathogenic Escherichia coli (APEC) infection. *PLoS One* **7**, e41645.

Ordas, A., Hegedus, Z., Henkel, C.V., Stockhammer, O.W., Butler, D., Jansen, H.J., *et al.* (2011). Deep sequencing of the innate immune transcriptomic response of zebrafish embryos to Salmonella infection. *Fish Shellfish Immun* **31**, 716-724.

Orskov, I., Svanborg Eden, C. and Orskov, F. (1988). Aerobactin production of serotyped Escherichia coli from urinary tract infections. *Medical microbiology and immunology* **177**, 9-14.

Pello, O.M., De Pizzol, M., Mirolo, M., Soucek, L., Zammataro, L., Amabile, A., *et al.* (2012). Role of c-MYC in alternative activation of human macrophages and tumor-associated macrophage biology. *Blood* **119**, 411-421.

Pichon, C., Hechard, C., du Merle, L., Chaudray, C., Bonne, I., Guadagnini, S., *et al.* (2009). Uropathogenic Escherichia coli AL511 requires flagellum to enter renal collecting duct cells. *Cell Microbiol* **11**, 616-628.

Plos, K., Carter, T., Hull, S., Hull, R. and Svanborg Eden, C. (1990). Frequency and organization of pap homologous DNA in relation to clinical origin of uropathogenic Escherichia coli. *The Journal of infectious diseases* **161**, 518-524.

- Plos, K., Connell, H., Jodal, U., Marklund, B.I., Marild, S., Wettergren, B. and Svanborg, C. (1995). Intestinal carriage of P fimbriated *Escherichia coli* and the susceptibility to urinary tract infection in young children. *J Infect Dis* **171**, 625-631.
- Pompella, A., Visvikis, A., Paolicchi, A., De Tata, V. and Casini, A.F. (2003). The changing faces of glutathione, a cellular protagonist. *Biochemical pharmacology* **66**, 1499-1503.
- Portt, L., Norman, G., Clapp, C., Greenwood, M. and Greenwood, M.T. (2011). Anti-apoptosis and cell survival: a review. *Biochimica et biophysica acta* **1813**, 238-259.
- Roberts, A., Trapnell, C., Donaghey, J., Rinn, J.L. and Pachter, L. (2011). Improving RNA-Seq expression estimates by correcting for fragment bias. *Genome Biol* **12**, R22.
- Roos, V., Nielsen, E.M. and Klemm, P. (2006). Asymptomatic bacteriuria *Escherichia coli* strains: adhesins, growth and competition. *FEMS Microbiol Lett* **262**, 22-30.
- Rosenberger, C.M. and Finlay, B.B. (2003). Phagocyte sabotage: disruption of macrophage signalling by bacterial pathogens. *Nature reviews. Molecular cell biology* **4**, 385-396.
- Rossol, M., Heine, H., Meusch, U., Quandt, D., Klein, C., Sweet, M.J. and Hauschildt, S. (2011). LPS-induced cytokine production in human monocytes and macrophages. *Critical reviews in immunology* **31**, 379-446.
- Salvatore, S., Salvatore, S., Cattoni, E., Siesto, G., Serati, M., Sorice, P. and Torella, M. (2011). Urinary tract infections in women. *Eur J Obstet Gynecol Reprod Biol* **156**, 131-136.
- Schembri, M.A., Sokurenko, E.V. and Klemm, P. (2000). Functional flexibility of the FimH adhesin: insights from a random mutant library. *Infection and immunity* **68**, 2638-2646.
- Semeraro, F., Ammollo, C.T., Morrissey, J.H., Dale, G.L., Friese, P., Esmon, N.L. and Esmon, C.T. (2011). Extracellular histones promote thrombin generation through platelet-dependent mechanisms: involvement of platelet TLR2 and TLR4. *Blood* **118**, 1952-1961.
- Sester, D.P., Beasley, S.J., Sweet, M.J., Fowles, L.F., Cronau, S.L., Stacey, K.J. and Hume, D.A. (1999). Bacterial/CpG DNA down-modulates colony stimulating factor-1 receptor surface expression on murine bone marrow-derived macrophages with concomitant growth arrest and factor-independent survival. *Journal of immunology* **163**, 6541-6550.
- Sester, D.P., Brion, K., Trieu, A., Goodridge, H.S., Roberts, T.L., Dunn, J., et al. (2006). CpG DNA activates survival in murine macrophages through TLR9 and the phosphatidylinositol 3-kinase-Akt pathway. *Journal of immunology* **177**, 4473-4480.
- Shih, A.Y., Erb, H., Sun, X., Toda, S., Kalivas, P.W. and Murphy, T.H. (2006). Cystine/glutamate exchange modulates glutathione supply for neuroprotection from oxidative stress and cell proliferation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **26**, 10514-10523.
- Sivick, K.E. and Mobley, H.L. (2010). Waging war against uropathogenic *Escherichia coli*: winning back the urinary tract. *Infect Immun* **78**, 568-585.
- Stenqvist, K., Sandberg, T., Lidin-Janson, G., Orskov, F., Orskov, I. and Svanborg-Eden, C. (1987). Virulence factors of *Escherichia coli* in urinary isolates from pregnant women. *The Journal of infectious diseases* **156**, 870-877.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

t Hoen, P.A., Ariyurek, Y., Thygesen, H.H., Vreugdenhil, E., Vossen, R.H., de Menezes, R.X., *et al.* (2008). Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms. *Nucleic Acids Res* **36**, e141.

Tan, C.K., Carey, A.J., Cui, X., Webb, R.I., Ipe, D., Crowley, M., *et al.* (2012). Genome-wide mapping of cystitis due to *Streptococcus agalactiae* and *Escherichia coli* in mice identifies a unique bladder transcriptome that signifies pathogen-specific antimicrobial defense against urinary tract infection. *Infection and immunity* **80**, 3145-3160.

Tegner, J., Nilsson, R., Bajic, V.B., Bjorkegren, J. and Ravasi, T. (2006). Systems biology of innate immunity. *Cell Immunol* **244**, 105-109.

Thomas-Chollier, M., Defrance, M., Medina-Rivera, A., Sand, O., Herrmann, C., Thieffry, D. and van Helden, J. RSAT 2011: regulatory sequence analysis tools. *Nucleic Acids Res* **39**, W86-91.

Thomas-Chollier, M., Sand, O., Turatsinze, J.V., Janky, R., Defrance, M., Vervisch, E., *et al.* (2008). RSAT: regulatory sequence analysis tools. *Nucleic Acids Res* **36**, W119-127.

Tomich, M., Herfst, C.A., Golden, J.W. and Mohr, C.D. (2002). Role of flagella in host cell invasion by *Burkholderia cepacia*. *Infection and immunity* **70**, 1799-1806.

Totsika, M., Moriel, D.G., Idris, A., Rogers, B.A., Wurpel, D.J., Phan, M.D., *et al.* (2012). Uropathogenic *Escherichia coli* mediated urinary tract infection. *Current drug targets* **13**, 1386-1399.

Tourneur, E., Ben Mkaddem, S., Chassin, C., Bens, M., Goujon, J.M., Charles, N., *et al.* (2013). Cyclosporine A impairs nucleotide binding oligomerization domain (Nod1)-mediated innate antibacterial renal defenses in mice and human transplant recipients. *PLoS pathogens* **9**, e1003152.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., *et al.* (2012). Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-578.

Turatsinze, J.V., Thomas-Chollier, M., Defrance, M. and van Helden, J. (2008). Using RSAT to scan genome sequences for transcription factor binding sites and cis-regulatory modules. *Nat Protoc* **3**, 1578-1588.

Ulett, G.C., Totsika, M., Schaale, K., Carey, A.J., Sweet, M.J. and Schembri, M.A. (2013). Uropathogenic *Escherichia coli* virulence and innate immune responses during urinary tract infection. *Curr Opin Microbiol* **16**, 100-107.

Wang, F., Hu, S., Liu, W., Qiao, Z., Gao, Y. and Bu, Z. (2012). Deep-sequencing analysis of the mouse transcriptome response to infection with *Brucella melitensis* strains of differing virulence. *PLoS One* **6**, e28485.

Wang, Z., Gerstein, M. and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* **10**, 57-63.

Welch, R.A., Burland, V., Plunkett, G., 3rd, Redford, P., Roesch, P., Rasko, D., *et al.* (2002). Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc Natl Acad Sci U S A* **99**, 17020-17024.

- Wiles, T.J., Kulesus, R.R. and Mulvey, M.A. (2008). Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol* **85**, 11-19.
- Willner, D., Low, S., Steen, J.A., George, N., Nimmo, G.R., Schembri, M.A. and Hugenholtz, P. (2014). Single clinical isolates from acute uncomplicated urinary tract infections are representative of dominant in situ populations. *mBio* **5**, e01064-01013.
- Xiang, L.X., He, D., Dong, W.R., Zhang, Y.W. and Shao, J.Z. (2010a). Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC Genomics* **11**.
- Xiang, L.X., He, D., Dong, W.R., Zhang, Y.W. and Shao, J.Z. (2010b). Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC Genomics* **11**, 472.
- Xiao, S., Jia, J., Mo, D., Wang, Q., Qin, L., He, Z., *et al.* (2010). Understanding PRRSV infection in porcine lung based on genome-wide transcriptome response identified by deep sequencing. *PLoS One* **5**, e11377.
- Xu, J., Zhang, X., Monestier, M., Esmon, N.L. and Esmon, C.T. (2011). Extracellular histones are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. *Journal of immunology* **187**, 2626-2631.
- Xu, J., Zhang, X., Pelayo, R., Monestier, M., Ammollo, C.T., Semeraro, F., *et al.* (2009). Extracellular histones are major mediators of death in sepsis. *Nature medicine* **15**, 1318-1321.
- Yamaguchi, S. and Darwin, A.J. (2012). Recent findings about the *Yersinia enterocolitica* phage shock protein response. *Journal of microbiology* **50**, 1-7.
- Yamaguchi, S., Reid, D.A., Rothenberg, E. and Darwin, A.J. (2013). Changes in Psp protein binding partners, localization and behaviour upon activation of the *Yersinia enterocolitica* phage shock protein response. *Mol Microbiol* **87**, 656-671.
- Zdziarski, J., Brzuszkiewicz, E., Wullt, B., Liesegang, H., Biran, D., Voigt, B., *et al.* (2010). Host imprints on bacterial genomes--rapid, divergent evolution in individual patients. *PLoS Pathog* **6**, e1001078.
- Zheng, M., Wang, X., Templeton, L.J., Smulski, D.R., LaRossa, R.A. and Storz, G. (2001). DNA microarray-mediated transcriptional profiling of the *Escherichia coli* response to hydrogen peroxide. *Journal of bacteriology* **183**, 4562-4570.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Figure Legends**

**Figure 1: BMM Transcriptome Analysis**

**A.** MCE plot of the relationships between conditions for the BMM gene sets. Each sample from two independent biological replicates is represented as a dot in a two-dimensional space (C: Control; U: UTI89; A: 83972). **B.** Venn diagrams quantifying the overlap in the response of BMMs to the two UPEC strains for two independent biological replicates. The numbers of DEG are shown for the total response (top left), as well as for each of the three time points (2, 4, and 24 hpi). **C.** Histogram of the regulation of BMM DEG showing the numbers of up- (red) and downregulated (green) genes during the 24-hour infection time course. **D.** Pathways activated in BMMs during the course of UPEC infection. Each colored square in the matrix represents a significant fold-enrichment ( $\log_2$ ) of the respective pathway term at each point. Red, upregulated DEG; green, downregulated DEG.

**Figure 2: Gene Regulation in BMM**

**A.** Heat map summarizing the expression profiles of histone genes from the RNA-Seq libraries. The values are log-transformed FPKM counts. **B.** Bar plots showing the relative mRNA levels of selected histone candidate genes determined by RT-qPCR. qPCR data represent means relative expression  $\pm$  range (n = 2 independent experiments). **C.** Bar plot showing the mean relative levels of the mRNA for Slc7a11, as determined by RT-qPCR. Error bars denote the range of the two biological replicates (C: Control; U: UTI89; A: 83972).

**Figure 3: Transcription Factors Associated with DE BMM Genes**

**A.** Heat map showing results of k-means clustering of BMM DEG. The values are log-transformed FPKM counts for all DEG across all RNA-Seq libraries in the dataset. **B.** Expression profiles of TFs

associated with binding motifs from the TFBS analysis that are highly correlated with each cluster. Each TF is represented with a different color, whereas the cluster mean expression is colored blue. All values are log-transformed FPKM counts. C: Control, U: UTI89, A: 83972

#### Figure 4: UPEC Transcriptome Analysis

**A.** MCE plot of the relationships between conditions for the UPEC gene sets. Each sample from two independent biological replicates is represented as a dot in a two-dimensional space (C: Control; U: UTI89; A: 83972). **B.** Venn diagrams quantifying the overlap in the response of the UPEC strains in the intramacrophage environment for two independent biological replicates. The numbers of DEG for the total response (top left), as well as for each of the three time points (2, 4, and 24 hpi), are shown. **C.** Histogram of the regulation of UPEC DEG showing the numbers of up- (red) and downregulated (green) genes during the 24-hour infection time. **D.** Pathways activated in UPEC during the course of infection. Each colored square in the matrix represents significant fold-enrichment ( $\log_2$ ) of the respective pathway term at each point. Red, upregulated DEG; green, downregulated DEG.

#### Figure 5: Regulation of UPEC Flagella, OxyR Regulon, and Hydrogen Peroxide-Induced Genes in the intramacrophage environment

**A-C.** Heat maps summarizing the RNA-Seq-derived expression profiles of flagellar genes (A), OxyR regulon genes (B), and hydrogen peroxide-induced genes (C). All values are log-transformed FPKM counts. **D.** Bar plot showing the relative mRNA levels of *ahpF*, as determined by RT-qPCR. qPCR data represent mean relative expression  $\pm$  range ( $n = 2$ ) of two biological replicates; ND: not detected.

#### Figure 6: UPEC Genes Associated with Intramacrophage Survival

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**A** and **B**. Heat maps summarizing the RNA-Seq-derived expression profiles of UTI89 genes elevated at 24 hpi (A) and UPEC Psp genes (B). **C**. Bar plots showing the relative quantity of *pspA* and *pspE* mRNA, as determined by RT-qPCR. qPCR data represent mean relative expression  $\pm$  range (n = 2) of two biological replicates; ND: not detected. **D**. Insertion site for creation of UTI89 *pspA* mutant. **E**. Intramacrophage survival of UTI89 and UTI89*pspA*. BMMs were infected at an MOI of 10 and intracellular bacterial survival was assessed at 1, 2 and 24 hours of infection. Data are compiled from three independent experiments, and show mean  $\pm$  standard deviation (\* p<0.05).

For Peer Review



## Supporting Information Captions

### **Supplementary Figure 1:** Intramacrophage survival of UTI89 versus 83972

Bacterial loads of UTI89 and 83972 within BMM at 2, 4 and 24 hpi in gentamicin exclusion assays were assessed by colony counting. These samples were used for RNA-Seq analyses. Data represent average cfu ml<sup>-1</sup> ± range (n = 2 independent experiments).

### **Supplementary Figure 2:** Enriched mouse and UPEC GO terms during the course of infection

Gene ontology terms enriched in BMMs (A) and UPEC (B) during the 24h course of infection. Each colored square in the matrix represents a significant fold-enrichment (log<sub>2</sub>) of the respective GO term at each point. Red, upregulated DEG; green, downregulated DEG.

### **Supplementary Figure 3:** Bioinformatic analysis pipeline

Summary of the steps followed for the generation and analysis of the RNA-Seq data produced by next-generation sequencing. Steps are grouped into 6 tiers, and details are provided on the algorithms, databases and software used for each of the analyses.

### **Supplementary Table 1:** Quality control of RNA-Seq libraries

### **Supplementary Table 2:** Alignment statistics of RNA-Seq reads

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**The Co-Transcriptome of Uropathogenic *Escherichia coli*-Infected Mouse Macrophages  
Reveals New Insights into Host–Pathogen Interactions**

Charalampos (Harris) Mavromatis<sup>1,2</sup>, Nilesh J. Bokil<sup>3,4</sup>, Makrina Totsika<sup>4,5,6</sup>, Asha Kakkanat<sup>4,5</sup>, Kolja  
Schaale<sup>3,4</sup>, Carlo V. Cannistraci<sup>1,2,7</sup>, Taewoo Ryu<sup>1,2</sup>, Scott A. Beatson<sup>4,5</sup>, Glen C Ulett<sup>8</sup>, Mark A.  
Schembri<sup>4,5</sup>, Matthew J. Sweet<sup>3,4</sup> and Timothy Ravasi<sup>1,2§</sup>

1. Division of Biological and Environmental Sciences and Engineering, Division of Computer, Electrical and Mathematical Sciences and Engineering, King Abdullah University of Science and Technology, Thuwal, Kingdom of Saudi Arabia
2. Division of Medical Genetics, Department of Medicine, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA USA
3. Institute for Molecular Bioscience, The University of Queensland, St Lucia, Australia
4. Australian Infectious Diseases Research Centre, The University of Queensland, St Lucia, Australia
5. School of Chemistry and Molecular Biosciences, The University of Queensland, St Lucia, Australia
6. Institute of Health and Biomedical Innovation (IHBI), School of Biomedical Sciences, Queensland University of Technology (QUT), 60 Musk Ave/cnr. Blamey St, Kelvin Grove, QLD, 4059, Australia
7. Biomedical Cybernetics Group, Biotechnology Center (BIOTEC), Technische Universität Dresden, Tatzberg 47/49, 01307 Dresden, Germany.
8. Griffith Health Institute and School of Medical Science, Griffith Health Centre, Gold Coast Campus, Griffith University, Southport, QLD 4222, Australia.

§ Corresponding author

**Email Addresses of Authors**

CM <harris.mavromatis@kaust.edu.sa>

NJB <n.bokil@imb.uq.edu.au>

MT <makrina.totsika@qut.edu.au>

AK <asha.kakkanat@uq.net.au>

KS <k.schaale@imb.uq.edu.au>

CVC <carlo.cannistraci@biotec.tu-dresden.de>

TR <taewoo.ryu@kaust.edu.sa>

SAB <s.beatson@uq.edu.au>

GCC <g.ulett@griffith.edu.au>

MAS <m.schembri@uq.edu.au>

MJS <m.sweet@imb.uq.edu.au>

TR <timothy.ravasi@kaust.edu.sa>

**Word Count: 6,735**

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Summary

Urinary tract infections (UTI) are among the most common infections in humans. Uropathogenic *Escherichia coli* (UPEC) can invade and replicate within bladder epithelial cells, and some UPEC strains can also survive within macrophages. To understand the UPEC transcriptional program associated with intramacrophage survival, we performed host–pathogen co-transcriptome analyses using RNA sequencing. Mouse bone marrow-derived macrophages (BMMs) were challenged over a 24 h time course with two UPEC reference strains that possess contrasting intramacrophage phenotypes: UTI89, which survives in BMMs, and 83972, which is killed by BMMs. Neither of these strains caused significant BMM cell death at the low multiplicity of infection that was used in this study. We developed an effective computational framework that simultaneously separated, annotated, and quantified the mammalian and bacterial transcriptomes. BMMs responded to the two UPEC strains with a broadly similar gene expression program. In contrast, the transcriptional responses of the UPEC strains diverged markedly from each other. We identified UTI89 genes upregulated at 24 h post-infection, and hypothesized that some may contribute to intramacrophage survival. Indeed, we showed that deletion of one such gene (*pspA*) significantly reduced UTI89 survival within BMMs. Our study provides a technological framework for simultaneously capturing global changes at the transcriptional level in co-cultures, and has generated new insights into the mechanisms that UPEC use to persist within the intramacrophage environment.

## Introduction

Urinary tract infections (UTIs) represent one of the most significant community-acquired and healthcare-associated diseases (Foxman, 2010, Horvath *et al.*, 2012). Uncomplicated UTIs result in more than 14 million medical visits and account for almost \$4 billion in medical expenditure each year in the USA alone (Salvatore *et al.*, 2011). Approximately 50% of women will experience a UTI at some point in their life, with almost 25% of patients experiencing a recurrence within the first 6 months following treatment of the initial UTI (Salvatore *et al.*, 2011). An estimated 68% of recurrent UTIs arise from the same bacterial strain that caused the initial infection (Hunstad *et al.*, 2010). Uropathogenic *Escherichia coli* (UPEC) is the most common causative agent of UTIs, being responsible for ~80% of all community-acquired infections (Foxman, 2010). In the majority of acute, uncomplicated UPEC-mediated UTIs, single cultured isolates are diagnostic of the infection (Willner *et al.*, 2014).

UPEC employ a range of virulence factors, including adhesins, toxins and iron-acquisition systems, to colonize the urinary tract and cause disease (Totsika *et al.*, 2012, Ulett *et al.*, 2013). Different UPEC strains display extensive genetic diversity owing to the presence of mobile DNA elements such as “pathogenicity islands”, prophages and plasmids (Hacker *et al.*, 2000, Mysorekar *et al.*, 2006, Wiles *et al.*, 2008, Hunstad *et al.*, 2010, Hannan *et al.*, 2012). The UPEC strains UTI89 (Mulvey *et al.*, 2001) and 83972 (Lindberg *et al.*, 1975c, Klemm *et al.*, 2007, Zdziarski *et al.*, 2010) are representative of cystitis and asymptomatic bacteriuria (ABU) isolates, respectively.

Acute pyelonephritis and ABU represent the two extremes of UTI. Acute pyelonephritis is a severe, acute systemic infection caused by UPEC strains containing virulence genes clustered on pathogenicity islands (Eden *et al.*, 1976, Funfstuck *et al.*, 1986, Stenqvist *et al.*, 1987, Orskov *et al.*, 1988, Johnson, 1991, Welch *et al.*, 2002). ABU, on the other hand, is an asymptomatic carrier state that resembles commensalism. A single *E. coli* strain may be present in ABU patients at levels of more

1  
2 than  $10^5$  colony-forming units (CFU)  $\text{ml}^{-1}$  for months or years without provoking a host response.  
3  
4 Because the majority of ABU-associated *E. coli* strains are non-hemolytic, non-adherent and lack  
5  
6 hemagglutination ability, early studies suggested that this behavior reflected a lack of virulence genes  
7  
8 (Lindberg, 1975, Lindberg *et al.*, 1975a, Lindberg *et al.*, 1975b, Lindberg *et al.*, 1975c, Eden *et al.*,  
9  
10 1976, Kaijser *et al.*, 1977). Molecular epidemiology has shown, however, that many ABU strains carry  
11  
12 virulence genes despite failing to express the phenotype (Plos *et al.*, 1990, Plos *et al.*, 1995, Mabbett  
13  
14 *et al.*, 2009).

15  
16  
17 As with all infectious agents, UPEC must overcome innate immunity, a biological system  
18  
19 compromising both cellular mediators (e.g. neutrophils) and soluble mediators (e.g. complement  
20  
21 proteins) that act synergistically. Several studies have investigated the role of neutrophils in UPEC-  
22  
23 mediated pathology (Ingersoll *et al.*, 2008, Sivick *et al.*, 2010, Lau *et al.*, 2012, Tourneur *et al.*, 2013),  
24  
25 whereas there is a paucity of information on the interactions between UPEC and macrophages,  
26  
27 another key cellular component of innate immunity (Tegner *et al.*, 2006). We previously demonstrated  
28  
29 that the ability of UPEC to survive in mouse macrophages differs markedly between different strains;  
30  
31 some strains, such as UTI89, are able to survive over a 24-hour infection period, whereas others, such  
32  
33 as 83972, are rapidly killed (Bokil *et al.*, 2011). This suggests that UPEC strains like UTI89 are able to  
34  
35 subvert macrophage antimicrobial pathways, though the mechanisms responsible are still unknown.  
36  
37 These findings are in keeping with a larger body of literature documenting intraepithelial cell survival of  
38  
39 some UPEC strains, both *in vitro* and *in vivo* (Hunstad *et al.*, 2010).  
40  
41  
42

43 Next-generation sequencing technologies provide a powerful approach for studying co-  
44  
45 transcriptomics during infection (t Hoen *et al.*, 2008, Hegedus *et al.*, 2009, Jager *et al.*, 2009, Xiang *et*  
46  
47 *al.*, 2010b, Huang *et al.*, 2012, Nie *et al.*, 2012, Wang *et al.*, 2012). These direct sequencing  
48  
49 methodologies allow the measurement of millions of RNA transcripts in a sample, thus enabling  
50  
51 identification of global differences in gene expression under different growth conditions (Morozova *et*  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 *al.*, 2008, Wang *et al.*, 2009). RNA sequencing generates information about absolute transcript levels,  
3  
4 avoiding many of the limitations of microarrays (t Hoen *et al.*, 2008, Llorens *et al.*, 2011). To further  
5  
6 understand the transcriptional programs that are simultaneously activated during host-pathogen  
7  
8 interaction, we developed an approach for isolating total RNA from co-cultures and analyzing the  
9  
10 simultaneous changes in expression that take place in the interacting organisms. Previous studies  
11  
12 (Hegedus *et al.*, 2009, Xiang *et al.*, 2010a, Xiao *et al.*, 2010, Ordas *et al.*, 2011) have focused on  
13  
14 either the host or the pathogen, without revealing simultaneous co-transcriptomic changes that occur  
15  
16 during infection. Such methods have also relied on the isolation of species-specific RNA, which can  
17  
18 introduce biases in the analysis.  
19  
20

21 To gain insights into novel strategies used by UPEC to subvert macrophage anti-microbial  
22  
23 responses, we performed a global co-transcriptomic analysis of UPEC gene expression within murine  
24  
25 bone marrow-derived macrophages (BMMs). We used RNA sequencing to monitor (i) the  
26  
27 transcriptional responses of UPEC strains UTI89 and 83972 within the intramacrophage environment  
28  
29 across an extended time course, and (ii) differences in macrophage gene-expression responses to  
30  
31 each strain. Our comprehensive approach has generated new insights into host–pathogen interactions  
32  
33 and the possible consequences of these interactions for disease processes.  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Results**

*Analysis of Digital Gene Expression Libraries*

The UPEC strains UTI89 and 83972 display contrasting intramacrophage survival phenotypes; UTI89 is able to survive in significant numbers, whereas 83972 is rapidly killed (Bokil *et al.*, 2011). To examine the molecular basis for this difference, we investigated the gene expression profiles of UTI89 and 83972 in BMMs in parallel over a 24 h period. For our experimental system, we used a multiplicity of infection (MOI) of 10:1, since this MOI does not have obvious effects on BMM cell viability for either of the strains. As expected, bacterial loads of UTI89 that were recovered from BMM were substantially higher than those of 83972 (Supplementary Figure 1). Total RNA was harvested at 2, 4 and 24 h post-infection (hpi) and global gene expression profiles were analyzed using the Illumina Hi-Seq 2000 Digital Gene Expression Tag Profiling Kit, a tag-based transcriptome sequencing method. cDNA libraries were prepared from gentamicin-treated UPEC-BMM co-cultures, sequenced and analyzed together with bacterial and BMM control samples. The RNA-Seq files generated for all libraries were preprocessed by a custom java script (Supplementary Table 1).

*Mapping RNA-Seq Reads*

Alignment of sequencing reads to the respective mouse and UPEC reference genomes was performed using TopHat. Sequences were mapped against their respective *Mus musculus* or *E. coli* genomes. The resulting BAM files were further used to compute alignment statistics for all libraries employing the samtool flagstat command. These processes are summarized in Supplementary Table 2. As expected, very few sequence reads were captured from 83972-BMM co-cultures at 24 hpi, consistent with the observation that these bacteria were essentially cleared by BMMs at this time point. Hence, subsequent analyses of gene expression in 83972 excluded this specific condition.



### BMM Transcriptional Responses to UPEC

To explore transcriptional relationships between different conditions, we performed dimensionality reduction analysis using minimum curvilinearity embedding (MCE) (Cannistraci *et al.*, 2010, Cannistraci *et al.*, 2013). This analysis revealed that the BMM genes whose expression was regulated by infection were not markedly different between the two different UPEC strains at the initial stage, 2hpi. However, after 4hpi, a slight difference emerged, which was also maintained at 24hpi. As expected, however, we observed that the time-dependent regulation of gene expression (Figure 1A) was the main pattern that emerged from the data, indicating that the macrophage transcriptional response follows a distinct temporal profile that is common to infection by both strains, and much more distinctive than the difference between the two strains.

To gain further insights into the global transcriptional changes that take place in UPEC-infected macrophages, we catalogued differentially expressed genes (DEG) in normalized digital gene expression data through pairwise comparisons between controls (uninfected BMM cultures) and treatments (co-cultures of UPEC-infected BMMs) using a previously described method (Trapnell *et al.*, 2012) with a threshold of a false-discovery rate (FDR)-adjusted *P*-value < 0.01 in at least one of the pairwise comparisons. Using this approach, we identified 628 and 652 BMM genes that were differentially regulated following infection with UTI89 or 83972, respectively, over a 24-hour infection time course. Further analyses of the BMM transcriptome identified 603 genes that were commonly regulated by UTI89 and 83972, as well as 25 and 49 BMM genes that were differentially regulated after infection with UTI89 or 83972, respectively (Figure 1B). The greatest divergence in responses to each strain occurred at the latest time point post-infection.

To investigate the regulatory patterns of divergently expressed (DE) genes, we clustered genes as either up or downregulated. This analysis revealed a rapid response for upregulated genes

1  
2 that was maintained throughout the infection time course. In contrast, the number of downregulated  
3  
4 genes was much lower at 2 and 4 hpi, but increased markedly by 24 hpi (Figure 1C). The substantial  
5  
6 overlap in DEG, as well as their conserved pattern of regulation, suggests that the macrophage  
7  
8 transcriptional response to UTI89 and 83972 is broadly conserved.  
9

10  
11  
12  
13 *Functional Annotation of DE Macrophage Genes*  
14

15 The consequences of gene expression changes associated with UPEC infection were  
16  
17 characterized by gene ontology (GO) and pathway (KEGG) enrichment analyses of DEG using the  
18  
19 DAVID program (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a). As shown in Supplementary Figure  
20  
21 2A, common highly enriched GO categories for upregulated genes included activation of inflammatory  
22  
23 responses; regulation of chemokine, cytokine, and interleukin-6 and -12 production; T-cell activation;  
24  
25 and regulation of transcription factor (TF) activity. Common biological processes associated with  
26  
27 downregulated genes included DNA-replication initiation, DNA packaging, nucleosome assembly and  
28  
29 organization, and cytokinesis. These pathways are consistent with the known activation of innate  
30  
31 immune responses by bacterial challenge (Rosenberger *et al.*, 2003, Mogensen, 2009, Portt *et al.*,  
32  
33 2011). As expected, the signaling pathways that were inferred, on the basis of gene expression  
34  
35 changes, to be regulated in macrophages upon UPEC infection showed common characteristics  
36  
37 between both UPEC strains. Signaling pathways associated with upregulated genes included  
38  
39 cytokine–cytokine receptor interaction, NOD-like receptor signaling, and Toll-like receptor (TLR)  
40  
41 signaling pathways (Figure 1D).  
42  
43  
44

45 We next independently validated both conservation and divergence in macrophage responses  
46  
47 to UTI89 versus 83972. Many of the well-validated TLR target genes such as cytokines (Il1a, Il1b, Il16)  
48  
49 and chemokines (Cxcl1, Ccl8) were similarly inducible by both UTI89 and 83972 in BMM (data not  
50  
51 shown). Although previous studies have demonstrated pathological roles for extracellular histones in  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 mouse models of sepsis (Xu *et al.*, 2009, Xu *et al.*, 2011), there is little known about the regulation of  
2 histone gene expression downstream of TLR signaling. This may reflect the fact that canonical histone  
3 mRNAs are not poly-adenylated, and so their regulated expression may not be captured by traditional  
4 microarray approaches. Interestingly, we found that a large suite of histone genes were  
5 downregulated in response to infection with either UPEC strain (Figure 2A), consistent with the  
6 inhibitory effect of TLR signaling on macrophage proliferation. This observation was validated for  
7 several individual histone genes using quantitative reverse transcription-polymerase chain reaction  
8 (RT-qPCR; Figure 2B). Our attempts to validate the small number of macrophage genes differentially  
9 regulated by UTI89 and 83972 (Figure 1B) using qPCR were less successful, however we did confirm  
10 differential regulation of the cystine/glutamate exchanger *Slc7a11*. Whereas *Slc7a11* mRNA  
11 expression was similarly upregulated by UTI89 and 83972 at 4 hpi in infected BMM, its expression  
12 remained elevated at 24 hpi in UTI89-infected BMMs but was significantly reduced at this time point in  
13 83972-infected BMMs (Figure 2C).

14  
15 Clustering analysis separated DE macrophage genes into two major clusters (Figure 3A).  
16 Cluster 1 (Figure 3B), which contained 460 genes, was positively correlated with the profiles of the  
17 TFs *Arnt* (aryl hydrocarbon receptor nuclear translocator), *Myc* and *Pparg* (peroxisome proliferator-  
18 activated receptor gamma); and negatively correlated with the profiles of *Hif1a* (hypoxia-inducible  
19 factor 1-alpha) and *Stat3* (signal transducer and activator of transcription 3); and enriched for  
20 pathways associated with DNA replication, cell cycle, and systemic lupus erythematosus. Cluster 2  
21 (Figure 3B), which contained 217 genes, was positively correlated with the profiles of the TFs *Hif1a*,  
22 *Nfkb2* (nuclear factor of kappa light polypeptide gene enhancer in B cells 2), *Nr3c1* (nuclear receptor  
23 subfamily 3, group C, member 1) and *Stat3*; negatively correlated with the profile of *Myc*; and enriched  
24 for pathways associated with cytokine–cytokine receptor interaction, and NOD-like receptor, TLR,  
25 chemokine and Jak-STAT signaling. These results are in keeping with expectations; TLR signaling

1  
2 activates **NF- $\kappa$ B** and **HIF** (Rossol *et al.*, 2011) and inactivates CSF-1 signaling. Various studies link  
3  
4 the biology of **CSF-1**, **Myc** and **Pparg** (Dey *et al.*, 2000, Bonfield *et al.*, 2008, Pello *et al.*, 2012), and as  
5  
6 expected, downregulated genes have an association with **Myc** and **Pparg**. That is, TLR signaling  
7  
8 switches off signaling via CSF-1, which itself can signal, in part, by **Myc**; hence, TLR signaling  
9  
10 downregulates **Myc** responses. Apart from the expected patterns, our analysis suggests a potential  
11  
12 association between inactivation of CSF-1 signaling and **Arnt** (partner for **Hif**) responses, which has  
13  
14 not been reported previously.  
15  
16  
17  
18

19 *UPEC Transcriptional Responses upon Infection of BMMs*

20  
21 As was the case with the macrophage gene expression analysis, we initially investigated  
22  
23 transcriptional responses in both UPEC strains by performing dimensionality reduction analysis using  
24  
25 MCE (Cannistraci *et al.*, 2010, Cannistraci *et al.*, 2013). We identified specific differences in the  
26  
27 regulated expression of UPEC genes within the intramacrophage environment (Figure 4A), which  
28  
29 likely reflects the differential pathogenicity and capacity for intramacrophage survival of both strains  
30  
31 (Supplementary Figure 1) (Bokil *et al.*, 2011).  
32  
33  
34

35 To gain insights into the global transcriptional changes that occur in UPEC during macrophage  
36  
37 infection, we applied the same method described above and performed pairwise comparisons  
38  
39 between controls (UPEC cultures alone) and treatments (co-cultures of UPEC-infected BMMs). Again,  
40  
41 we employed an FDR-adjusted *P*-value < 0.01 in at least one of the pairwise comparisons in our  
42  
43 dataset as the threshold for differential expression. In total, we identified 137 UTI89 genes and 33  
44  
45 83972 genes that were differentially regulated in BMMs. Surprisingly, an analysis of the  
46  
47 intramacrophage transcriptomic profile identified only 13 regulated genes that were common to both  
48  
49 strains, whereas 124 genes were uniquely regulated in UTI89, and 20 genes were uniquely regulated  
50  
51 in 83972 (Figure 4B). The UPEC transcriptional profiles also revealed a number of strain-specific  
52  
53  
54  
55  
56  
57  
58  
59  
60

1 responses; a substantial number of UTI89 genes were upregulated at 2 and 4 hpi, whereas a much  
2 smaller number of 83972 genes were upregulated at these time points (Figure 4C). This is consistent  
3 with the initiation of a UTI89 transcriptional program that permits intramacrophage survival. In contrast  
4 to the transcriptional responses observed for macrophage genes, few UPEC genes were  
5 downregulated over the infection time course.  
6  
7  
8  
9  
10  
11  
12  
13  
14

#### 15 *Functional Annotation of DE UPEC Genes*

16  
17 The consequences of gene expression changes associated with UPEC infection were  
18 characterized by GO and pathway (KEGG) enrichment analyses of DEG using the DAVID program  
19 (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a). This again provided clear evidence of strain-specific  
20 biological responses within the intramacrophage environment (Supplementary Figure 2B). Highly  
21 enriched GO categories associated with the upregulated transcriptional program included genes  
22 associated with chemotaxis and motility in UTI89 and genes involved in protein folding in 83972. We  
23 did not identify any significantly enriched GO categories associated with downregulated genes in  
24 common for the two strains. In keeping with the above findings, an analysis of the pathways activated  
25 during UTI89 infection revealed enrichment for pathway terms associated with bacterial chemotaxis  
26 and flagellar biosynthesis (Figure 4D). We did not observe significant enrichment for any pathway term  
27 in the case of 83972.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42

#### 43 *UPEC Genes Associated with Intra-Macrophage Survival*

44  
45 We hypothesised that genes selectively upregulated by UTI89 may contribute to  
46 intramacrophage survival. Such genes included those encoding flagella and those associated with  
47 protection against oxidative stress. Notably, several genes encoding flagella-related proteins were  
48 uniquely regulated in UTI89; these DE flagella genes included *flgA-F*, *flgK*, *flgL*, *fliC-E*, *fliQ*, *motA*, and  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 *motB*. Figure 5A shows the expression patterns of all flagella-related genes in UTI89 versus 83972,  
3  
4 revealing strong upregulation at 2 hours with a subsequent gradual decrease over time.

5  
6 A common mechanism used by bacterial pathogens to avoid host innate immune pathways is  
7  
8 to employ defense mechanisms against oxidative stress (Imlay, 2013). We therefore clustered the  
9  
10 expression patterns of all the OxyR regulon genes (Figure 5B) and the most strongly hydrogen  
11  
12 peroxide-induced genes in both UPEC strains (Figure 5C). Interestingly, the OxyR regulon, which  
13  
14 included alkyl hydroperoxide reductase subunit F (*ahpF*), *dps* (DNA starvation/stationary phase  
15  
16 protection protein), *grxA* (glutaredoxin 1), *trxC* (thioredoxin 2) and *yaaA*, was strongly upregulated in  
17  
18 UTI89, with the effect being apparent at 2 hpi and peaking at 4 hpi. Hydrogen peroxide-inducible  
19  
20 genes, which included *ahpF*, *dps*, *grxA*, heat-shock proteins/chaperones (*ibpA*, *ibpB*), *phoH*, *soxS*  
21  
22 (DNA-binding transcriptional dual regulator), *trxC* and *yaaA* showed a similar pattern, with genes being  
23  
24 significantly upregulated in UTI89 and less so in 83972. Differential regulation of *ahpF* in UTI89 versus  
25  
26 83972 within macrophages was validated by RT-qPCR (Figure 5D).  
27  
28  
29

30 Although UTI89 persists at 24 h within macrophages, only a relatively small percentage of  
31  
32 bacteria (< 5%) survive within BMM at 24 hpi compared to the bacterial loads at 2 hpi (Supplementary  
33  
34 Figure 1). We therefore reasoned that changes in UPEC gene expression at 24 hpi might be linked to  
35  
36 intramacrophage survival. We identified 22 genes that were highly upregulated (> 3-fold) by UTI89 at  
37  
38 24 hpi (Figure 6A). The most highly expressed of these included those encoding *ibpB* (encoding a  
39  
40 small heat shock protein); *pspACDE* (encoding the phage-shock protein system); *rpoE* and *rpoH*  
41  
42 (encoding sigma factors); *smpA* (encoding outer membrane lipoprotein); *yadR* (encoding the iron-  
43  
44 sulfur cluster insertion protein ErpA); *yceP* (encoding the biofilm formation regulatory protein BssS);  
45  
46 *yebG* (encoding DNA damage-inducible protein); and UTI89\_C2624 and UTI89\_C5162-3 (encoding  
47  
48 proteins of unknown function).  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 Phage-shock-protein (Psp)-related genes, which are required for bacterial survival during  
3  
4 extracytoplasmic stress responses and changes in pH (Darwin, 2013), were significantly upregulated  
5  
6 in UTI89 compared to 83972 (Figure 6B). RT-qPCR confirmed the elevated expression of *pspA* and  
7  
8 *pspE* in UTI89 compared to 83972 at 24 h post-infection (Figure 6C). Finally, we validated the impact  
9  
10 of *pspA* for intramacrophage survival of UTI89 by constructing a UTI89*pspA* mutant (Figure 6D), and  
11  
12 testing it for intramacrophage survival during a 24 h infection time course in BMM. In this assay, the  
13  
14 UTI89*pspA* mutant was significantly reduced for intracellular survival compared to the wild type strain  
15  
16 (Figure 6E).  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Discussion**

In this study, we determined the co-transcriptomic program of UPEC-infected primary mouse macrophages during an infection time course. The use of two UPEC strains that differed in their ability to survive in these cells enabled the identification of both common and UPEC strain-specific responses. To our knowledge, this is one of the first RNA-Seq studies that have simultaneously measured the transcriptomes of both the host and pathogen during an infection (Humphrys *et al.*, 2013). Using only open-source tools, we developed a computational framework that was capable of successfully separating, annotating, and quantifying the mammalian and bacterial transcriptomes. Whereas previous studies (Mysorekar *et al.*, 2002, Bower *et al.*, 2009, Hagan *et al.*, 2010, Duell *et al.*, 2012) have been limited to microarrays, the RNA-Seq analysis reported here provides a more sensitive, comprehensive, and unbiased coverage of the entire transcriptome. We achieved an average sequencing depth of approximately 24 million tags per library and identified 677 BMM and 157 UPEC genes that were differentially expressed following UPEC infection. By mapping our RNA-Seq tag data onto transcript databases and genomic sequences, we were able to identify genes that were regulated upon UPEC challenge.

MCE confirmed the distinct temporal cascade of BMM responses to UPEC as well as differences in transcriptional programs of the two UPEC strains. The sets of DEG and regulatory processes and pathways identified in both BMMs and UPEC strains suggest coordinated expression and mutual influences. Our study detected 603 DEG in BMMs that were common to infection with UTI89 and 83972, accounting for the vast majority of changes in BMM gene expression. This was not unexpected, given that both UTI89 and 83972 present pathogen-associated molecular patterns, such as lipopolysaccharide that is recognized by TLR4. Macrophage-expressed genes that showed differences in regulation in response to these two strains were a very minor component of the total



signature and their significance remains to be determined. Despite the general conservation of the BMM response against UPEC infection, we nonetheless expected to detect some selective BMM responses at 24 hpi for UTI89 compared with 83972, since the former condition reflects macrophages that are continuing to cope with intracellular UPEC, whereas the latter corresponds to macrophages that have cleared the infection. Consistent with this supposition, we found that at 24 hpi, BMM responses showed a greater divergence between UTI89 and 83972 (47.8% overlap) compared to earlier time points where this difference was not as evident. Of interest in this regard, was our validation of differential regulation of the cationic amino acid cysteine/glutamate antiporter Slc7a11, which is required for glutamate uptake and glutathione synthesis (Bannai, 1986, Hayes *et al.*, 1999, Pompella *et al.*, 2003, Shih *et al.*, 2006). This gene showed a sustained upregulation at 24 hpi following infection with UTI89, whereas it was only transiently upregulated after infection with 83972. Given that macrophage antimicrobial responses typically involve subjecting pathogens to oxidative stress, the sustained expression of Slc7a11 upon infection with UTI89 may be required to maintain glutathione levels and cytoprotection during stress responses.

Our data is consistent with previous reports showing inducible cytokine and chemokine expression, as well as activation of pro-survival pathways (Sester *et al.*, 1999, Sester *et al.*, 2006), during bacterial infection and/or TLR stimulation of macrophages. Moreover, the downregulation of DNA replication and cell cycle genes (Figure 1D) is consistent with the known inhibitory effects of TLR4 signalling on proliferation of cycling macrophages (Sester *et al.*, 1999, Sester *et al.*, 2006). We also found that the genes encoding histones H1, H2, and H4 were dramatically downregulated at 24 hpi, and we further confirmed the regulated expression of several of these using RT-qPCR. A member of the H2A histone family, *Hist2h2aa1*, was also significantly downregulated at 24 hpi *in vivo* in mouse bladder colonized with UPEC CFT073 (Tan *et al.*, 2012), which is consistent with these findings. Interestingly, previous studies have reported a pathological role for extracellular histones during LPS-

1 induced septic shock (Xu *et al.*, 2009, Li *et al.*, 2011). Xu *et al.* (2011) revealed that antibodies against  
2 extracellular histones rescued animals from LPS-mediated death (Xu *et al.*, 2011), and a previous  
3 study showed that extracellular histones mediate endothelial dysfunction, organ failure, and death  
4 during sepsis (Semeraro *et al.*, 2011). Our findings could thus either reflect a host attempt to reduce  
5 inflammatory responses upon cell death and histone release, or the consequence of growth-inhibitory  
6 effects of TLR agonists on proliferating macrophages, which would also be expected to lead to  
7 downregulated histone expression.  
8  
9  
10  
11  
12  
13  
14  
15  
16

17 A transcription factor binding site (TFBS) analysis of the promoter sequences of the two BMM  
18 DE gene clusters further showed significant enrichment of seven motifs associated with TFs that have  
19 key roles in macrophage functions. The cluster containing **inducible** inflammation-related genes  
20 correlated positively with the expression of **Hif1a**, a key pro-inflammatory transcription factor that  
21 drives macrophage inflammatory responses and is upregulated in UPEC-infected mouse bladder at 2  
22 hpi and 24 hpi (Duell *et al.*, 2012, Tan *et al.*, 2012); **Nfkb2**, which is activated by a wide variety of  
23 stimuli such as cytokines, oxidant-free radicals and bacterial or viral products; the glucocorticoid  
24 receptor **Nr3c1**, which up-regulates the expression of anti-inflammatory genes and/or represses the  
25 expression of pro-inflammatory genes; and **Stat3**, a transcriptional activator stimulated in response to  
26 cytokines and growth factors. **In vivo**, both **Nfkb2** and **Stat3** are immediately upregulated in the bladder  
27 following UPEC infection (Duell *et al.*, 2012). On the other hand, the cluster containing **downregulated**  
28 cell cycle-related genes correlated positively with the expression of **Arnt**, which is involved in the  
29 induction of several enzymes that participate in xenobiotic metabolism; **Myc**, which is **required for cell**  
30 **proliferation in response to mitogenic stimuli such as CSF-1**; and **Pparg**, which regulates fatty acid  
31 storage and glucose metabolism. **Myc**, **Pparg**, and **Csf1** are all differentially expressed in bladders of  
32 mice infected with UPEC (Duell *et al.*, 2012) supporting the *in vivo* relevance of these findings.  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 In contrast to the general conserved pattern of BMM responses to both UPEC strains, the  
3  
4 individual transcriptional responses of the two strains in BMMs differed markedly. This is consistent  
5  
6 with their contrasting survival patterns in macrophages. MCE confirmed the different transcriptional  
7  
8 programs of the two UPEC strains, revealing 124 UTI89-specific DEG, 20 83972-specific DEG, and 13  
9  
10 DEG that were commonly regulated in both pathogens upon infection of BMMs. Given the different  
11  
12 survival patterns of both UPEC strains in macrophages, we expected to detect increased expression  
13  
14 of genes specific to UTI89 within macrophages at 24 hpi.  
15  
16

17 Previous studies have shown that flagella contribute to the virulence of a number of pathogenic  
18  
19 species (Tomich *et al.*, 2002). Some bacteria, for example *Salmonella enterica* Typhimurium and  
20  
21 *Yersinia enterocolitica*, use flagella for invasion of epithelial cells (McNally *et al.*, 2007, Ibarra *et al.*,  
22  
23 2010). UPEC also employ flagella for the invasion of renal collecting duct cells (Pichon *et al.*, 2009),  
24  
25 but flagella-enhanced uptake of UPEC by macrophages has not been investigated. Our co-  
26  
27 transcriptomic analysis revealed the upregulation of multiple genes associated with flagella  
28  
29 biosynthesis. This could reflect inducible gene expression in the intramacrophage environment or  
30  
31 selective uptake by macrophages of a sub-population of flagella-expressing UPEC, which would  
32  
33 generate a similar gene expression profile. The acute upregulation at 2 hpi and subsequent  
34  
35 downregulation at 24 hpi would suggest the latter may be the case.  
36  
37

38 Murine macrophages employ both rapidly produced reactive oxygen species (ROS) and  
39  
40 reactive nitrogen species (RNS), produced in a delayed fashion, as bacterial clearance strategies  
41  
42 (Flannagan *et al.*, 2009). The observation that UPEC strain UTI89 survives in macrophages suggests  
43  
44 that it can overcome these pathways. Although the genes that respond to ROS have been well  
45  
46 characterized in *E. coli* K-12, their expression during UPEC infection and their role in intramacrophage  
47  
48 survival have not yet been studied. Our transcriptomic approach identified UPEC genes that are likely  
49  
50 involved in defense against ROS and RNS. Disturbances in the normal redox state of cells can cause  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 toxic effects through the production of peroxides and free radicals that damage all components of the  
3  
4 cell, including proteins, lipids, and DNA. An earlier study confirmed that the peroxide response  
5  
6 regulator OxyR activates most of the genes that are highly induced by hydrogen peroxide (Zheng *et*  
7  
8 *al.*, 2001). Our results show that the response of UTI89 to oxidative stress (upregulated expression of  
9  
10 *dps*, *grxA*, *ahpF*, *soxS*, *trxC*, *ibpA*, and *ibpB*) is more robust than that of 83972. These differences  
11  
12 imply that part of the UTI89 intramacrophage survival strategy includes robust protection against  
13  
14 oxidative stress.  
15  
16

17  
18 Several UTI89 genes were upregulated in BMM at 24 hpi, and we selected genes encoding the  
19  
20 Psp system for further functional analysis. The Psp system responds to extracytoplasmic stress and  
21  
22 contributes to the virulence of several pathogens, including *Y. enterocolitica* and *S. Typhimurium*  
23  
24 (Karlinsey *et al.*, 2010, Yamaguchi *et al.*, 2012). PspA, the master effector of the Psp system,  
25  
26 mediates its response via a dual mechanism: (i) binding to the transcriptional regulator PspF and  
27  
28 preventing it from activating the transcription of *pspACDE* in the absence of extracytoplasmic stress,  
29  
30 and (ii) binding to the cytoplasmic membrane-localized proteins PspB and PspC in the presence of  
31  
32 extracytoplasmic stress, thus releasing PspF to induce *pspACDE* transcription (Yamaguchi *et al.*,  
33  
34 2013). We confirmed a role for the Psp system in UPEC by constructing a *pspA* mutant (UTI89*pspA*)  
35  
36 and demonstrating reduced intramacrophage survival of this mutant. Despite this statistically  
37  
38 significant reduction in survival within macrophages, the biological significance of this effect is at this  
39  
40 stage unknown. Inactivation of other stress response systems on the *pspA* mutant background may be  
41  
42 required to reveal a more striking phenotype.  
43  
44

45  
46 In summary, we have demonstrated the capacity to employ co-transcriptomics to study host-  
47  
48 pathogen interactions. Our novel approach revealed new insights into the mechanisms used by UPEC  
49  
50 to avoid macrophage responses and persist in the intra-macrophage environment, and has identified  
51  
52 multiple target genes for further functional studies. Finally, this manuscript is one of the first to  
53  
54  
55  
56  
57  
58  
59  
60

1  
2 successfully evaluate the expression profiles of two organisms from the same sample using RNA  
3  
4 sequencing, and will make an excellent resource for other studies that aim to perform similar analyses  
5  
6 in different host-pathogen systems.  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Peer Review

1  
2 **Experimental Procedures**

3  
4  
5  
6 *TIER I: In vitro Infection Assays*

7  
8 Ethics Statement and Animal Experimentation

9  
10 A University of Queensland institutional animal ethics committee approved all animal  
11 experimentations. Female C57BL/6 mice (6–8 weeks old) were purchased from the Animal Resources  
12 Center, Australia. Murine bone marrow-derived macrophages (BMM) were generated by the *in vitro*  
13 differentiation of bone marrow cells from C57BL/6 mice on bacteriological plastic plates in the  
14 presence of 10,000 U ml<sup>-1</sup> recombinant human CSF-1 (a gift from Chiron) for 6 days, after which cells  
15 were harvested and replated onto tissue culture plastic in the presence of CSF-1 for infection on day  
16 7. BMM were maintained at 37°C (5% CO<sub>2</sub>) in RPMI 1640 supplemented with 2mM L-glutamine  
17 (GlutaMAX), 10% heat-inactivated fetal bovine serum and 50 U ml<sup>-1</sup> penicillin and 50 µg ml<sup>-1</sup>  
18 streptomycin (Life Technologies).  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31

32 Culture of Bacterial Strains and Infection of Mouse BMMs

33  
34 *E. coli* UTI89 is a well-characterized cystitis isolate (Mulvey *et al.*, 2001). *E. coli* 83972, which  
35 was carried without symptoms by a young female, was originally isolated from the urine of this  
36 individual (Andersson *et al.*, 1991, Roos *et al.*, 2006). For macrophage infection assays, UPEC strains  
37 were cultured statically in Luria-Bertani (LB) broth at 37°C overnight. Type 1 fimbriae expression was  
38 assessed by yeast cell agglutination prior to infection as previously described (Schembri *et al.*, 2000).  
39 Bacterial cells were centrifuged and washed in phosphate buffered saline (PBS), and then  
40 resuspended in antibiotic-free media at a concentration of 2 x 10<sup>8</sup> CFU ml<sup>-1</sup>. Viable CFU counts of  
41 bacterial inocula were routinely confirmed in every infection assay by serial dilution and plating on LB  
42 agar.  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Intramacrophage survival assays were performed essentially as described previously (Bokil *et al.*, 2011). Briefly, following overnight adherence in antibiotic-free media, BMMs were infected for 1h with bacteria at an MOI of 10. Extracellular bacteria were killed by washing twice in 200  $\mu\text{g ml}^{-1}$  gentamicin, followed by 1h incubation in media containing the same gentamicin concentration. Subsequent exclusion of extracellular bacteria for the duration of the experiment was performed by incubation in 20  $\mu\text{g ml}^{-1}$  gentamicin. At appropriate time points (1h, 2h and 24h), cells were washed twice with antibiotic-free media, and then lysed with PBS/0.01% Triton X-100. Lysates were cultured on LB agar plates overnight at 37°C and colony counts were used to assess intra-cellular bacterial loads. For standard infection assays, complete exclusion of viable extracellular bacteria was confirmed by performing colony counts on culture supernatants.

## *TIER II: RNA Preparation and Sequencing*

### Macrophage Infection and mRNA Isolation, Enrichment, and Purification

Control macrophages, control bacteria and infected macrophages were incubated for an additional 1, 3 and 23 h at 37°C in a humidified 5% CO<sub>2</sub> atmosphere, after the initial 1 h infection. Cells were then washed twice with antibiotic-free media, and then lysed on ice for RNA isolation and purification (RNeasy; Qiagen, Germantown, MD, USA). Microbial total RNA in co-culture samples was enriched (MICROBEnrich, Ambion). rRNA was removed from all purified RNA samples using kits targeting mammalian and Gram-negative bacterial rRNAs (Ribo-Zero; Epicenter, Madison, WI, USA). Prior to sequencing, all samples were further quantified and examined for protein and reagent contamination using a Nanodrop ND-1000 spectrophotometer. RNA samples for analysis were selected based on a 28S/18S rRNA band intensity of 2:1, a spectroscopic  $A_{260}/A_{280}$  nm ratio of 1.8–2.0, and an  $A_{260}/A_{230}$  nm ratio > 1.5.



1  
2 RNA Sequencing

3  
4       Next-generation sequencing analyses were performed for two biological replicates on an  
5  
6 Illumina Cluster Station and the Illumina HiSeq 2000 System using primarily reagents from the Illumina  
7  
8 Gene Expression Sample Preparation Kit and the Illumina Sequencing Chip (Flowcell; Illumina, San  
9  
10 Diego, CA, USA). Sequence tags were prepared using the Digital Gene Expression Tag Profiling Kit  
11  
12 (Illumina), according to the Illumina protocol.  
13  
14

15  
16  
17 *TIER III: Data Pre-processing*

18  
19       Image analysis, base calling, and quality calibration were performed using the Solexa  
20  
21 Automated Pipeline. Quality control of RNA-Seq reads were preprocessed by a custom java script. All  
22  
23 sequences generated have a length of 101 bases and have been submitted to the BioProject  
24  
25 database of NCBI under BioProject ID: PRJNA256028.  
26  
27

28  
29  
30 *TIER IV: Alignment and Differential Gene Expression Analysis*

31  
32 Alignment of Reads

33  
34       Bowtie indexes were created for the mouse (version 37.1) and the two *E. coli* strains (UTI89  
35  
36 and 83972) using the bowtie-build algorithm and reference sequences from the GenBank database.  
37  
38 Our protocol began with mapping of the raw RNA-Seq reads (fastq files) to the reference genomes  
39  
40 using TopHat. TopHat uses Bowtie as an alignment engine and breaks up reads that Bowtie cannot  
41  
42 align on its own into smaller pieces (Kim *et al.*, 2011). Using the standard parameters, we mapped  
43  
44 both reads of our paired-end libraries. All simulations were performed using a 30-core, high-memory  
45  
46 node cluster system; total computation duration was 3 hours.  
47  
48

49  
50  
51 Transcript Annotation

After running TopHat, the resulting alignment files were provided to Cufflinks to generate a transcriptome assembly for each condition. During this analysis step, adapter tags; mitochondrial sequences; poly A, poly C, and phiX sequences; and remaining ribosomal sequences were filtered out. Estimated normalized expression levels were reported in Fragments (i.e., reads) Per Kilobase of exon per Million mapped reads (FPKM). These assemblies were compared with annotation files using the Cuffcompare utility, which is included in the Cufflinks package (Roberts *et al.*, 2011, Trapnell *et al.*, 2012).

### Differential Expression Analysis

The reads and assemblies were imported to Cuffdiff, which calculates expression levels and tests the statistical significance of observed changes. For comparison of DEG across samples, the number of raw clean tags in each library was normalized to FPKM using the Cufflinks package. The minimum number of alignments in a locus needed to test for significance of changes in that locus between samples was set to 50 fragment alignments. If no testing was performed, changes in the locus were deemed insignificant, and the changes observed in the locus did not contribute to corrections for multiple testing. The Cuffdiff output files were then imported to cummeRbund, which plots abundance and differential expression results as commonly used expression plots for quality control (Trapnell *et al.*, 2012).

### *TIER V: Functional Analysis*

#### Dimensionality Reduction

Dimensionality reduction is necessary for exploring the relationships between conditions in our experiment. The MCE method performs a nonlinear dimension reduction by embedding high-dimensional data points into a lower-dimensional space using the minimum curvilinear kernel in

1 combination with multidimensional scaling (MDS) (Cannistraci *et al.*, 2010) or in alternative the  
2  
3 singular value decomposition (Cannistraci *et al.*, 2013). The nonlinear data distances for MDS or SVD  
4  
5 were computed and stored in the minimum curvilinear kernel as the traversal distances over the  
6  
7 minimum spanning tree between the data points (in our study the samples' conditions) in the  
8  
9 multidimensional space (in our case the gene space). The minimum spanning tree was constructed  
10  
11 from the Pearson correlation-based distances between the samples (Cannistraci *et al.*, 2010):  
12  
13

14  
15 
$$\text{correlation\_based\_distance}(x,y) = 1 - \text{Pearson\_correlation}(\text{Sample}_x, \text{Sample}_y)$$
  
16

17 MCE is a parameter-free projection algorithm that was shown to be particularly effective in  
18  
19 discriminating classes in small-*n* (samples: here conditions), large-*m* (features: here gene  
20  
21 expressions) datasets using only the first dimension of embedding (Cannistraci *et al.*, 2010). The fact  
22  
23 that our datasets have  $n \ll m$  led us to adopt MCE algorithm for unsupervised analysis of the patterns  
24  
25 present between the different sample conditions.  
26  
27

28  
29  
30 Clustering of Genes  
31

32 Genes with similar expression patterns often serve overlapping functions. Accordingly, the  
33  
34 optimal number of clusters in the dataset was determined by performing a cluster analysis of gene  
35  
36 expression patterns using the R package NbClust. Selected lists of expression profiles of DE genes  
37  
38 were compiled for each hypothesis tested and clustered using the Ward's methodology.  
39  
40  
41

42  
43 GO and Pathway-enrichment analyses of DEG  
44

45 Genes involved in common biological processes or pathways tend to show overlapping  
46  
47 expression profiles. In gene expression profiling analyses, significantly enriched GO terms and  
48  
49 pathways were identified by mapping all DEG to terms in the GO and KEGG databases by applying  
50  
51 two-sided Fisher's exact and  $\chi^2$  tests, respectively (Huang da *et al.*, 2009b, Huang da *et al.*, 2009a).  
52  
53  
54  
55  
56  
57  
58  
59  
60

P-values were corrected by calculating the FDR, and only GO and pathway terms with a FDR < 0.01 were chosen.

### Identification of key BMM TFs

Promoter sequences of all DEG were retrieved using the RSAT and were further input into the RSAT matrix-scan tool along with mouse-related JASPAR matrices for TFBS prediction (Thomas-Chollier *et al.*, Thomas-Chollier *et al.*, 2008, Turatsinze *et al.*, 2008). The RSAT output was filtered using an adjusted P-value < 0.05 as a cut-off, and lists of the most significant TFBSs and their known corresponding TFs were compiled. The expression profiles of the mouse DEG were clustered and each cluster was correlated with TFs profiles using Pearson correlation in R. Finally, the clusters were annotated using GO to fully elucidate the molecular processes in which each TF was involved.

### Screening for strain-specific UPEC gene expression patterns

Flagella-related gene lists were compiled based on Macnab's review on bacterial flagellar assembly (Macnab, 2003). A list of 35 genes related to the flagellar apparatus was used as background for screening our UPEC datasets. For OxyR regulon and hydrogen peroxide-induced genes, gene lists were similarly compiled using previously published data on the response of bacteria to hydrogen peroxide (Zheng *et al.*, 2001). Finally, lists of genes involved in Psp regulation were compiled, and our datasets were screened for their expression patterns. The expression patterns of the screened genes were further clustered as described above, and all results were visualized using pheatmap package in R.

### Identification of UPEC Genes Associated with Intramacrophage Survival

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Putative bacterial genes associated with intramacrophage survival were considered as those that remained highly upregulated at 24 hpi. Bacterial genes upregulated at this time point were compiled and filtered based on their significance of expression. All DEG were filtered using an FDR-adjusted P-value of the test statistic  $< 0.01$  and a  $\log_2$  fold change  $> 3$ . The filtered lists of DEG enabled us to screen for survival genes, cluster their expression, and further annotate them using the GO database to better understand the biological processes they regulated.

*TIER VI: Gene Validation*

cDNA synthesis, primer design and RT-qPCR

cDNA was synthesized using a SuperScript III First-Strand Synthesis kit (Invitrogen). qPCR was then performed on a 7900HT Fast Real-Time PCR system (Applied Biosystems) using RNA samples from two independent biological replicates, similar to those employed for the RNA-Seq experiments. TaqMan Fast Universal PCR Master Mix 2X (Applied Biosystems) was used for BMM gene validation, and SYBR Green Master Mix (Applied Biosystems) was used for UPEC gene validation. Each cDNA was analyzed in triplicate, after which the average threshold cycle (Ct) per sample was calculated. Raw data were processed with qBase Plus software (Biogazelle), which performs downstream processing of qPCR data. The geNorm algorithm, integrated in the qBase Plus package, was used for determining the optimal number and identity of reference genes needed to normalize the data in both BMM (Actb and Polr2a) and UPEC (gapA and purC) qPCR libraries. Relative expression levels were calculated with the  $2^{-\Delta\Delta Ct}$  method ( $\Delta Ct$  is the difference in Ct between the reference genes and the target gene products); the average Ct value for all genes was used to correct for differences in cDNA input. Other statistical procedures were performed with the R program. All steps, from the experimental design to bioinformatic analysis and gene validation, are summarized in Supplementary Figure 3.

### Construction of UTI89 $\psi$ spA deletion mutant

Chromosomal DNA purification, PCR and DNA sequencing of PCR products was performed as previously described (Allsopp *et al.*, 2010). The  $\psi$ spA gene was mutated in UTI89 using the  $\lambda$ -Red mediated homologous recombination method with some modifications (Datsenko *et al.*, 2000, Allsopp *et al.*, 2012). Briefly, a three-step PCR procedure was employed to generate a DNA fragment comprising the chloramphenicol cassette from plasmid pKD3 and two 500-bp fragments homologous to the flanking regions of the  $\psi$ spA gene. The following primers were used: 5376\_UTI89 $\psi$ spA FwUP (5'-gccgtagcgagttcatca) and 5377\_UTI89 $\psi$ spA RvUP (5'-ggaataggaactaaggaggaagcgttgatgttggcatt), 5378\_UTI89 $\psi$ spA Fwdn (5'-cctacacaatcgctcaagacgccgaactgaaagccgat) and 5379\_UTI89 $\psi$ spA Rvsn (5'-taaacagcgccagaccga) to generate the 500bp homology arms; 3746-Cm.3a (5'-tcctccttagttcctattcc) and 3747-Cm.4a (5'-gtcttgagcgattgtgtagg) to generate the chloramphenicol resistance gene fragment. This DNA fusion product was electroporated into UTI89 harboring plasmid pKD46, and chloramphenicol resistant mutants were selected and confirmed by PCR (using primers 5375\_UTI89 $\psi$ spA FwSc: 5'-tcgtcgcgcataccaacc and 5380\_UTI89 $\psi$ spA Rvsc: 5'-acttcatccagcaattcgc). The UTI89 $\psi$ spA mutant was confirmed by sequencing.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Acknowledgments**

This work was supported by grants from the National Health and Medical Research Council (NHMRC) of Australia (APP1005315 and APP1068593). MJS is supported by an Australian Research Council (ARC) Future Fellowship (FT100100657), as well as an honorary NHMRC Senior Research Fellowship (APP1003470). MAS and GCU are supported by ARC Future Fellowships (FT100100662 and FT110101048, respectively), MT is supported by an ARC Discovery Early Career Researcher Award (DE130101169), and SAB is supported by an NHMRC Career Development Fellowship (APP1090456).



## References

- Allsopp, L.P., Beloin, C., Ulett, G.C., Valle, J., Totsika, M., Sherlock, O., *et al.* (2012). Molecular characterization of UpaB and UpaC, two new autotransporter proteins of uropathogenic *Escherichia coli* CFT073. *Infection and immunity* **80**, 321-332.
- Allsopp, L.P., Totsika, M., Tree, J.J., Ulett, G.C., Mabbett, A.N., Wells, T.J., *et al.* (2010). UpaH is a newly identified autotransporter protein that contributes to biofilm formation and bladder colonization by uropathogenic *Escherichia coli* CFT073. *Infection and immunity* **78**, 1659-1669.
- Andersson, P., Engberg, I., Lidin-Janson, G., Lincoln, K., Hull, R., Hull, S. and Svanborg, C. (1991). Persistence of *Escherichia coli* bacteriuria is not determined by bacterial adherence. *Infection and immunity* **59**, 2915-2921.
- Bannai, S. (1986). Exchange of cystine and glutamate across plasma membrane of human fibroblasts. *J Biol Chem* **261**, 2256-2263.
- Bokil, N.J., Totsika, M., Carey, A.J., Stacey, K.J., Hancock, V., Saunders, B.M., *et al.* (2011). Intramacrophage survival of uropathogenic *Escherichia coli*: differences between diverse clinical isolates and between mouse and human macrophages. *Immunobiology* **216**, 1164-1171.
- Bonfield, T.L., Thomassen, M.J., Farver, C.F., Abraham, S., Koloze, M.T., Zhang, X., *et al.* (2008). Peroxisome proliferator-activated receptor-gamma regulates the expression of alveolar macrophage colony-stimulating factor. *Journal of immunology* **181**, 235-242.
- Bower, J.M., Gordon-Raagas, H.B. and Mulvey, M.A. (2009). Conditioning of uropathogenic *Escherichia coli* for enhanced colonization of host. *Infect Immun* **77**, 2104-2112.
- Cannistraci, C.V., Alanis-Lobato, G. and Ravasi, T. (2013). Minimum curvilinearity to enhance topological prediction of protein interactions by network embedding. *Bioinformatics* **29**, i199-209.
- Cannistraci, C.V., Ravasi, T., Montevecchi, F.M., Ideker, T. and Alessio, M. (2010). Nonlinear dimension reduction and clustering by Minimum Curvilinearity unfold neuropathic pain and tissue embryological classes. *Bioinformatics* **26**, i531-539.
- Darwin, A.J. (2013). Stress relief during host infection: The phage shock protein response supports bacterial virulence in various ways. *PLoS pathogens* **9**, e1003388.
- Datsenko, K.A. and Wanner, B.L. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 6640-6645.
- Dey, A., She, H., Kim, L., Boruch, A., Guris, D.L., Carlberg, K., *et al.* (2000). Colony-stimulating factor-1 receptor utilizes multiple signaling pathways to induce cyclin D2 expression. *Molecular biology of the cell* **11**, 3835-3848.
- Duell, B.L., Carey, A.J., Tan, C.K., Cui, X., Webb, R.I., Totsika, M., *et al.* (2012). Innate transcriptional networks activated in bladder in response to uropathogenic *Escherichia coli* drive diverse biological pathways and rapid synthesis of IL-10 for defense against bacterial urinary tract infection. *Journal of immunology* **188**, 781-792.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Eden, C.S., Hanson, L.A., Jodal, U., Lindberg, U. and Akerlund, A.S. (1976). Variable adherence to normal human urinary-tract epithelial cells of *Escherichia coli* strains associated with various forms of urinary-tract infection. *Lancet* **1**, 490-492.

Flannagan, R.S., Cosio, G. and Grinstein, S. (2009). Antimicrobial mechanisms of phagocytes and bacterial evasion strategies. *Nature reviews. Microbiology* **7**, 355-366.

Foxman, B. (2010). The epidemiology of urinary tract infection. *Nat Rev Urol* **7**, 653-660.

Funfstuck, R., Tschape, H., Stein, G., Kunath, H., Bergner, M. and Wessel, G. (1986). Virulence properties of *Escherichia coli* strains in patients with chronic pyelonephritis. *Infection* **14**, 145-150.

Hacker, J. and Kaper, J.B. (2000). Pathogenicity islands and the evolution of microbes. *Annu Rev Microbiol* **54**, 641-679.

Hagan, E.C., Lloyd, A.L., Rasko, D.A., Faerber, G.J. and Mobley, H.L. (2010). *Escherichia coli* global gene expression in urine from women with urinary tract infection. *PLoS Pathog* **6**, e1001187.

Hannan, T.J., Totsika, M., Mansfield, K.J., Moore, K.H., Schembri, M.A. and Hultgren, S.J. (2012). Host-pathogen checkpoints and population bottlenecks in persistent and intracellular uropathogenic *Escherichia coli* bladder infection. *FEMS Microbiol Rev* **36**, 616-648.

Hayes, J.D. and McLellan, L.I. (1999). Glutathione and glutathione-dependent enzymes represent a co-ordinately regulated defence against oxidative stress. *Free radical research* **31**, 273-300.

Hegedus, Z., Zakrzewska, A., Agoston, V.C., Ordas, A., Racz, P., Mink, M., *et al.* (2009). Deep sequencing of the zebrafish transcriptome response to mycobacterium infection. *Mol Immunol* **46**, 2918-2930.

Horvath, D.J., Jr., Dabdoub, S.M., Li, B., Vanderbrink, B.A. and Justice, S.S. (2012). New paradigms of urinary tract infections: Implications for patient management. *Indian J Urol* **28**, 154-158.

Huang da, W., Sherman, B.T. and Lempicki, R.A. (2009a). Bioinformatics enrichment tools: paths toward the comprehensive functional analysis of large gene lists. *Nucleic Acids Res* **37**, 1-13.

Huang da, W., Sherman, B.T. and Lempicki, R.A. (2009b). Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* **4**, 44-57.

Huang, Q., Dong, S., Fang, C., Wu, X., Ye, T. and Lin, Y. (2012). Deep sequencing-based transcriptome profiling analysis of *Oryzias melastigma* exposed to PFOS. *Aquat Toxicol* **120-121**, 54-58.

Humphrys, M.S., Creasy, T., Sun, Y., Shetty, A.C., Chibucos, M.C., Drabek, E.F., *et al.* (2013). Simultaneous transcriptional profiling of bacteria and their host cells. *PLoS One* **8**, e80597.

Hunstad, D.A. and Justice, S.S. (2010). Intracellular lifestyles and immune evasion strategies of uropathogenic *Escherichia coli*. *Annu Rev Microbiol* **64**, 203-221.

- Ibarra, J.A., Knodler, L.A., Sturdevant, D.E., Virtaneva, K., Carmody, A.B., Fischer, E.R., *et al.* (2010). Induction of Salmonella pathogenicity island 1 under different growth conditions can affect Salmonella-host cell interactions in vitro. *Microbiology* **156**, 1120-1133.
- Imlay, J.A. (2013). The molecular mechanisms and physiological consequences of oxidative stress: lessons from a model bacterium. *Nature reviews. Microbiology* **11**, 443-454.
- Ingersoll, M.A., Kline, K.A., Nielsen, H.V. and Hultgren, S.J. (2008). G-CSF induction early in uropathogenic Escherichia coli infection of the urinary tract modulates host immunity. *Cellular microbiology* **10**, 2568-2578.
- Jager, D., Sharma, C.M., Thomsen, J., Ehlers, C., Vogel, J. and Schmitz, R.A. (2009). Deep sequencing analysis of the Methanosarcina mazei Go1 transcriptome in response to nitrogen availability. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 21878-21882.
- Johnson, J.R. (1991). Virulence factors in Escherichia coli urinary tract infection. *Clin Microbiol Rev* **4**, 80-128.
- Kaijser, B. and Ahlstedt, S. (1977). Protective capacity of antibodies against Escherichia coli and K antigens. *Infection and immunity* **17**, 286-289.
- Karlinsey, J.E., Maguire, M.E., Becker, L.A., Crouch, M.L. and Fang, F.C. (2010). The phage shock protein PspA facilitates divalent metal transport and is required for virulence of Salmonella enterica sv. Typhimurium. *Molecular microbiology* **78**, 669-685.
- Kim, D. and Salzberg, S.L. (2011). TopHat-Fusion: an algorithm for discovery of novel fusion transcripts. *Genome Biol* **12**, R72.
- Klemm, P., Hancock, V. and Schembri, M.A. (2007). Mellowing out: adaptation to commensalism by Escherichia coli asymptomatic bacteriuria strain 83972. *Infect Immun* **75**, 3688-3695.
- Lau, M.E., Loughman, J.A. and Hunstad, D.A. (2012). YbcL of uropathogenic Escherichia coli suppresses transepithelial neutrophil migration. *Infection and immunity* **80**, 4123-4132.
- Li, Y., Liu, B., Fukudome, E.Y., Lu, J., Chong, W., Jin, G., *et al.* (2011). Identification of citrullinated histone H3 as a potential serum protein biomarker in a lethal model of lipopolysaccharide-induced shock. *Surgery* **150**, 442-451.
- Lindberg, U. (1975). Asymptomatic bacteriuria in school girls. V. The clinical course and response to treatment. *Acta paediatrica Scandinavica* **64**, 718-724.
- Lindberg, U., Bjure, J., Haugstvedt, S. and Jodal, U. (1975a). Asymptomatic bacteriuria in schoolgirls. III. Relation between residual urine volume and recurrence. *Acta paediatrica Scandinavica* **64**, 437-440.
- Lindberg, U., Claesson, I., Hanson, L.A. and Jodal, U. (1975b). Asymptomatic bacteriuria in schoolgirls. I. Clinical and laboratory findings. *Acta paediatrica Scandinavica* **64**, 425-431.
- Lindberg, U., Hanson, L.A., Jodal, U., Lidin-Janson, G., Lincoln, K. and Olling, S. (1975c). Asymptomatic bacteriuria in schoolgirls. II. Differences in escherichia coli causing asymptomatic bacteriuria. *Acta paediatrica Scandinavica* **64**, 432-436.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Llorens, F., Hummel, M., Pastor, X., Ferrer, A., Pluvinet, R., Vivancos, A., *et al.* (2011). Multiple platform assessment of the EGF dependent transcriptome by microarray and deep tag sequencing analysis. *BMC Genomics* **12**, 326.

Mabbett, A.N., Ulett, G.C., Watts, R.E., Tree, J.J., Totsika, M., Ong, C.L., *et al.* (2009). Virulence properties of asymptomatic bacteriuria Escherichia coli. *International journal of medical microbiology : IJMM* **299**, 53-63.

Macnab, R.M. (2003). How bacteria assemble flagella. *Annu Rev Microbiol* **57**, 77-100.

McNally, A., La Ragione, R.M., Best, A., Manning, G. and Newell, D.G. (2007). An aflagellate mutant Yersinia enterocolitica biotype 1A strain displays altered invasion of epithelial cells, persistence in macrophages, and cytokine secretion profiles in vitro. *Microbiology* **153**, 1339-1349.

Mogensen, T.H. (2009). Pathogen recognition and inflammatory signaling in innate immune defenses. *Clinical microbiology reviews* **22**, 240-273, Table of Contents.

Morozova, O. and Marra, M.A. (2008). Applications of next-generation sequencing technologies in functional genomics. *Genomics* **92**, 255-264.

Mulvey, M.A., Schilling, J.D. and Hultgren, S.J. (2001). Establishment of a persistent Escherichia coli reservoir during the acute phase of a bladder infection. *Infection and immunity* **69**, 4572-4579.

Mysorekar, I.U. and Hultgren, S.J. (2006). Mechanisms of uropathogenic Escherichia coli persistence and eradication from the urinary tract. *Proc Natl Acad Sci U S A* **103**, 14170-14175.

Mysorekar, I.U., Mulvey, M.A., Hultgren, S.J. and Gordon, J.I. (2002). Molecular regulation of urothelial renewal and host defenses during infection with uropathogenic Escherichia coli. *J Biol Chem* **277**, 7412-7419.

Nie, Q., Sandford, E.E., Zhang, X., Nolan, L.K. and Lamont, S.J. (2012). Deep sequencing-based transcriptome analysis of chicken spleen in response to avian pathogenic Escherichia coli (APEC) infection. *PLoS One* **7**, e41645.

Ordas, A., Hegedus, Z., Henkel, C.V., Stockhammer, O.W., Butler, D., Jansen, H.J., *et al.* (2011). Deep sequencing of the innate immune transcriptomic response of zebrafish embryos to Salmonella infection. *Fish Shellfish Immun* **31**, 716-724.

Orskov, I., Svanborg Eden, C. and Orskov, F. (1988). Aerobactin production of serotyped Escherichia coli from urinary tract infections. *Medical microbiology and immunology* **177**, 9-14.

Pello, O.M., De Pizzol, M., Mirolo, M., Soucek, L., Zammataro, L., Amabile, A., *et al.* (2012). Role of c-MYC in alternative activation of human macrophages and tumor-associated macrophage biology. *Blood* **119**, 411-421.

Pichon, C., Hechard, C., du Merle, L., Chaudray, C., Bonne, I., Guadagnini, S., *et al.* (2009). Uropathogenic Escherichia coli AL511 requires flagellum to enter renal collecting duct cells. *Cell Microbiol* **11**, 616-628.

Plos, K., Carter, T., Hull, S., Hull, R. and Svanborg Eden, C. (1990). Frequency and organization of pap homologous DNA in relation to clinical origin of uropathogenic Escherichia coli. *The Journal of infectious diseases* **161**, 518-524.



- Plos, K., Connell, H., Jodal, U., Marklund, B.I., Marild, S., Wettergren, B. and Svanborg, C. (1995). Intestinal carriage of P fimbriated *Escherichia coli* and the susceptibility to urinary tract infection in young children. *J Infect Dis* **171**, 625-631.
- Pompella, A., Visvikis, A., Paolicchi, A., De Tata, V. and Casini, A.F. (2003). The changing faces of glutathione, a cellular protagonist. *Biochemical pharmacology* **66**, 1499-1503.
- Portt, L., Norman, G., Clapp, C., Greenwood, M. and Greenwood, M.T. (2011). Anti-apoptosis and cell survival: a review. *Biochimica et biophysica acta* **1813**, 238-259.
- Roberts, A., Trapnell, C., Donaghey, J., Rinn, J.L. and Pachter, L. (2011). Improving RNA-Seq expression estimates by correcting for fragment bias. *Genome Biol* **12**, R22.
- Roos, V., Nielsen, E.M. and Klemm, P. (2006). Asymptomatic bacteriuria *Escherichia coli* strains: adhesins, growth and competition. *FEMS Microbiol Lett* **262**, 22-30.
- Rosenberger, C.M. and Finlay, B.B. (2003). Phagocyte sabotage: disruption of macrophage signalling by bacterial pathogens. *Nature reviews. Molecular cell biology* **4**, 385-396.
- Rossol, M., Heine, H., Meusch, U., Quandt, D., Klein, C., Sweet, M.J. and Hauschildt, S. (2011). LPS-induced cytokine production in human monocytes and macrophages. *Critical reviews in immunology* **31**, 379-446.
- Salvatore, S., Salvatore, S., Cattoni, E., Siesto, G., Serati, M., Sorice, P. and Torella, M. (2011). Urinary tract infections in women. *Eur J Obstet Gynecol Reprod Biol* **156**, 131-136.
- Schembri, M.A., Sokurenko, E.V. and Klemm, P. (2000). Functional flexibility of the FimH adhesin: insights from a random mutant library. *Infection and immunity* **68**, 2638-2646.
- Semeraro, F., Ammollo, C.T., Morrissey, J.H., Dale, G.L., Friese, P., Esmon, N.L. and Esmon, C.T. (2011). Extracellular histones promote thrombin generation through platelet-dependent mechanisms: involvement of platelet TLR2 and TLR4. *Blood* **118**, 1952-1961.
- Sester, D.P., Beasley, S.J., Sweet, M.J., Fowles, L.F., Cronau, S.L., Stacey, K.J. and Hume, D.A. (1999). Bacterial/CpG DNA down-modulates colony stimulating factor-1 receptor surface expression on murine bone marrow-derived macrophages with concomitant growth arrest and factor-independent survival. *Journal of immunology* **163**, 6541-6550.
- Sester, D.P., Brion, K., Trieu, A., Goodridge, H.S., Roberts, T.L., Dunn, J., et al. (2006). CpG DNA activates survival in murine macrophages through TLR9 and the phosphatidylinositol 3-kinase-Akt pathway. *Journal of immunology* **177**, 4473-4480.
- Shih, A.Y., Erb, H., Sun, X., Toda, S., Kalivas, P.W. and Murphy, T.H. (2006). Cystine/glutamate exchange modulates glutathione supply for neuroprotection from oxidative stress and cell proliferation. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **26**, 10514-10523.
- Sivick, K.E. and Mobley, H.L. (2010). Waging war against uropathogenic *Escherichia coli*: winning back the urinary tract. *Infect Immun* **78**, 568-585.
- Stenqvist, K., Sandberg, T., Lidin-Janson, G., Orskov, F., Orskov, I. and Svanborg-Eden, C. (1987). Virulence factors of *Escherichia coli* in urinary isolates from pregnant women. *The Journal of infectious diseases* **156**, 870-877.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

t Hoen, P.A., Ariyurek, Y., Thygesen, H.H., Vreugdenhil, E., Vossen, R.H., de Menezes, R.X., *et al.* (2008). Deep sequencing-based expression analysis shows major advances in robustness, resolution and inter-lab portability over five microarray platforms. *Nucleic Acids Res* **36**, e141.

Tan, C.K., Carey, A.J., Cui, X., Webb, R.I., Ipe, D., Crowley, M., *et al.* (2012). Genome-wide mapping of cystitis due to *Streptococcus agalactiae* and *Escherichia coli* in mice identifies a unique bladder transcriptome that signifies pathogen-specific antimicrobial defense against urinary tract infection. *Infection and immunity* **80**, 3145-3160.

Tegner, J., Nilsson, R., Bajic, V.B., Bjorkegren, J. and Ravasi, T. (2006). Systems biology of innate immunity. *Cell Immunol* **244**, 105-109.

Thomas-Chollier, M., Defrance, M., Medina-Rivera, A., Sand, O., Herrmann, C., Thieffry, D. and van Helden, J. RSAT 2011: regulatory sequence analysis tools. *Nucleic Acids Res* **39**, W86-91.

Thomas-Chollier, M., Sand, O., Turatsinze, J.V., Janky, R., Defrance, M., Vervisch, E., *et al.* (2008). RSAT: regulatory sequence analysis tools. *Nucleic Acids Res* **36**, W119-127.

Tomich, M., Herfst, C.A., Golden, J.W. and Mohr, C.D. (2002). Role of flagella in host cell invasion by *Burkholderia cepacia*. *Infection and immunity* **70**, 1799-1806.

Totsika, M., Moriel, D.G., Idris, A., Rogers, B.A., Wurpel, D.J., Phan, M.D., *et al.* (2012). Uropathogenic *Escherichia coli* mediated urinary tract infection. *Current drug targets* **13**, 1386-1399.

Tourneur, E., Ben Mkaddem, S., Chassin, C., Bens, M., Goujon, J.M., Charles, N., *et al.* (2013). Cyclosporine A impairs nucleotide binding oligomerization domain (Nod1)-mediated innate antibacterial renal defenses in mice and human transplant recipients. *PLoS pathogens* **9**, e1003152.

Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D.R., *et al.* (2012). Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* **7**, 562-578.

Turatsinze, J.V., Thomas-Chollier, M., Defrance, M. and van Helden, J. (2008). Using RSAT to scan genome sequences for transcription factor binding sites and cis-regulatory modules. *Nat Protoc* **3**, 1578-1588.

Ulett, G.C., Totsika, M., Schaale, K., Carey, A.J., Sweet, M.J. and Schembri, M.A. (2013). Uropathogenic *Escherichia coli* virulence and innate immune responses during urinary tract infection. *Curr Opin Microbiol* **16**, 100-107.

Wang, F., Hu, S., Liu, W., Qiao, Z., Gao, Y. and Bu, Z. (2012). Deep-sequencing analysis of the mouse transcriptome response to infection with *Brucella melitensis* strains of differing virulence. *PLoS One* **6**, e28485.

Wang, Z., Gerstein, M. and Snyder, M. (2009). RNA-Seq: a revolutionary tool for transcriptomics. *Nat Rev Genet* **10**, 57-63.

Welch, R.A., Burland, V., Plunkett, G., 3rd, Redford, P., Roesch, P., Rasko, D., *et al.* (2002). Extensive mosaic structure revealed by the complete genome sequence of uropathogenic *Escherichia coli*. *Proc Natl Acad Sci U S A* **99**, 17020-17024.

- Wiles, T.J., Kulesus, R.R. and Mulvey, M.A. (2008). Origins and virulence mechanisms of uropathogenic *Escherichia coli*. *Exp Mol Pathol* **85**, 11-19.
- Willner, D., Low, S., Steen, J.A., George, N., Nimmo, G.R., Schembri, M.A. and Hugenholtz, P. (2014). Single clinical isolates from acute uncomplicated urinary tract infections are representative of dominant in situ populations. *mBio* **5**, e01064-01013.
- Xiang, L.X., He, D., Dong, W.R., Zhang, Y.W. and Shao, J.Z. (2010a). Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC Genomics* **11**.
- Xiang, L.X., He, D., Dong, W.R., Zhang, Y.W. and Shao, J.Z. (2010b). Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC Genomics* **11**, 472.
- Xiao, S., Jia, J., Mo, D., Wang, Q., Qin, L., He, Z., *et al.* (2010). Understanding PRRSV infection in porcine lung based on genome-wide transcriptome response identified by deep sequencing. *PLoS One* **5**, e11377.
- Xu, J., Zhang, X., Monestier, M., Esmon, N.L. and Esmon, C.T. (2011). Extracellular histones are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. *Journal of immunology* **187**, 2626-2631.
- Xu, J., Zhang, X., Pelayo, R., Monestier, M., Ammollo, C.T., Semeraro, F., *et al.* (2009). Extracellular histones are major mediators of death in sepsis. *Nature medicine* **15**, 1318-1321.
- Yamaguchi, S. and Darwin, A.J. (2012). Recent findings about the *Yersinia enterocolitica* phage shock protein response. *Journal of microbiology* **50**, 1-7.
- Yamaguchi, S., Reid, D.A., Rothenberg, E. and Darwin, A.J. (2013). Changes in Psp protein binding partners, localization and behaviour upon activation of the *Yersinia enterocolitica* phage shock protein response. *Mol Microbiol* **87**, 656-671.
- Zdziarski, J., Brzuszkiewicz, E., Wullt, B., Liesegang, H., Biran, D., Voigt, B., *et al.* (2010). Host imprints on bacterial genomes--rapid, divergent evolution in individual patients. *PLoS Pathog* **6**, e1001078.
- Zheng, M., Wang, X., Templeton, L.J., Smulski, D.R., LaRossa, R.A. and Storz, G. (2001). DNA microarray-mediated transcriptional profiling of the *Escherichia coli* response to hydrogen peroxide. *Journal of bacteriology* **183**, 4562-4570.

1  
2 **Figure Legends**

3  
4  
5  
6 **Figure 1: BMM Transcriptome Analysis**

7  
8 **A.** MCE plot of the relationships between conditions for the BMM gene sets. Each sample from two  
9 independent biological replicates is represented as a dot in a two-dimensional space (C: Control; U:  
10 UTI89; A: 83972). **B.** Venn diagrams quantifying the overlap in the response of BMMs to the two  
11 UPEC strains for two independent biological replicates. The numbers of DEG are shown for the total  
12 response (top left), as well as for each of the three time points (2, 4, and 24 hpi). **C.** Histogram of the  
13 regulation of BMM DEG showing the numbers of up- (red) and downregulated (green) genes during  
14 the 24-hour infection time course. **D.** Pathways activated in BMMs during the course of UPEC  
15 infection. Each colored square in the matrix represents a significant fold-enrichment ( $\log_2$ ) of the  
16 respective pathway term at each point. Red, upregulated DEG; green, downregulated DEG.

17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30 **Figure 2: Gene Regulation in BMM**

31  
32 **A.** Heat map summarizing the expression profiles of histone genes from the RNA-Seq libraries. The  
33 values are log-transformed FPKM counts. **B.** Bar plots showing the relative mRNA levels of selected  
34 histone candidate genes determined by RT-qPCR. qPCR data represent means relative expression  $\pm$   
35 range (n = 2 independent experiments). **C.** Bar plot showing the mean relative levels of the mRNA for  
36 Slc7a11, as determined by RT-qPCR. Error bars denote the range of the two biological replicates (C:  
37 Control; U: UTI89; A: 83972).

38  
39  
40  
41  
42  
43  
44  
45  
46  
47 **Figure 3: Transcription Factors Associated with DE BMM Genes**

48  
49 **A.** Heat map showing results of k-means clustering of BMM DEG. The values are log-transformed  
50 FPKM counts for all DEG across all RNA-Seq libraries in the dataset. **B.** Expression profiles of TFs



associated with binding motifs from the TFBS analysis that are highly correlated with each cluster. Each TF is represented with a different color, whereas the cluster mean expression is colored blue. All values are log-transformed FPKM counts. C: Control, U: UTI89, A: 83972

#### Figure 4: UPEC Transcriptome Analysis

**A.** MCE plot of the relationships between conditions for the UPEC gene sets. Each sample from two independent biological replicates is represented as a dot in a two-dimensional space (C: Control; U: UTI89; A: 83972). **B.** Venn diagrams quantifying the overlap in the response of the UPEC strains in the intramacrophage environment for two independent biological replicates. The numbers of DEG for the total response (top left), as well as for each of the three time points (2, 4, and 24 hpi), are shown. **C.** Histogram of the regulation of UPEC DEG showing the numbers of up- (red) and downregulated (green) genes during the 24-hour infection time. **D.** Pathways activated in UPEC during the course of infection. Each colored square in the matrix represents significant fold-enrichment ( $\log_2$ ) of the respective pathway term at each point. Red, upregulated DEG; green, downregulated DEG.

#### Figure 5: Regulation of UPEC Flagella, OxyR Regulon, and Hydrogen Peroxide-Induced Genes in the intramacrophage environment

**A-C.** Heat maps summarizing the RNA-Seq-derived expression profiles of flagellar genes (A), OxyR regulon genes (B), and hydrogen peroxide-induced genes (C). All values are log-transformed FPKM counts. **D.** Bar plot showing the relative mRNA levels of *ahpF*, as determined by RT-qPCR. qPCR data represent mean relative expression  $\pm$  range ( $n = 2$ ) of two biological replicates; ND: not detected.

#### Figure 6: UPEC Genes Associated with Intramacrophage Survival

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**A** and **B**. Heat maps summarizing the RNA-Seq-derived expression profiles of UTI89 genes elevated at 24 hpi (A) and UPEC Psp genes (B). **C**. Bar plots showing the relative quantity of *pspA* and *pspE* mRNA, as determined by RT-qPCR. qPCR data represent mean relative expression  $\pm$  range (n = 2) of two biological replicates; ND: not detected. **D**. Insertion site for creation of UTI89 *pspA* mutant. **E**. Intramacrophage survival of UTI89 and UTI89*pspA*. BMMs were infected at an MOI of 10 and intracellular bacterial survival was assessed at 1, 2 and 24 hours of infection. Data are compiled from three independent experiments, and show mean  $\pm$  standard deviation (\* p<0.05).

For Peer Review

## Supporting Information Captions

### **Supplementary Figure 1:** Intramacrophage survival of UTI89 versus 83972

Bacterial loads of UTI89 and 83972 within BMM at 2, 4 and 24 hpi in gentamicin exclusion assays were assessed by colony counting. These samples were used for RNA-Seq analyses. Data represent average cfu ml<sup>-1</sup> ± range (n = 2 independent experiments).

### **Supplementary Figure 2:** Enriched mouse and UPEC GO terms during the course of infection

Gene ontology terms enriched in BMMs (A) and UPEC (B) during the 24h course of infection. Each colored square in the matrix represents a significant fold-enrichment (log<sub>2</sub>) of the respective GO term at each point. Red, upregulated DEG; green, downregulated DEG.

### **Supplementary Figure 3:** Bioinformatic analysis pipeline

Summary of the steps followed for the generation and analysis of the RNA-Seq data produced by next-generation sequencing. Steps are grouped into 6 tiers, and details are provided on the algorithms, databases and software used for each of the analyses.

### **Supplementary Table 1:** Quality control of RNA-Seq libraries

### **Supplementary Table 2:** Alignment statistics of RNA-Seq reads




Figure 1  
183x122mm (300 x 300 DPI)




Figure 2  
137x74mm (300 x 300 DPI)




Figure 3  
152x106mm (300 x 300 DPI)




Figure 4  
160x93mm (300 x 300 DPI)




Figure 5  
101x59mm (300 x 300 DPI)






Figure 6  
157x94mm (300 x 300 DPI)